

HIGHER EULER-KRONECKER CONSTANTS

BY

SAMPRIT GHOSH

A thesis submitted in conformity with
the requirements for the degree of

Doctor of Philosophy

Department of Mathematics

University of Toronto

© 2023 Samprit Ghosh

ABSTRACT

Higher Euler-Kronecker Constants

Samprit Ghosh

Doctor of Philosophy

Department of Mathematics, University of Toronto

The coefficients that appear in the Laurent series of Dedekind zeta functions and their logarithmic derivatives are mysterious and seem to contain a lot of arithmetic information. Although the residue and the constant term have been widely studied, not much is known about the higher coefficients. In this thesis, we study these coefficients $\gamma_{K,n}$ that appear in the Laurent series expansion of $\frac{\zeta'_K(s)}{\zeta_K(s)}$ about $s = 1$, where K is a global field. For example, when K is a number field, we unconditionally prove certain arithmetic formulas satisfied by these coefficients and we give bounds for them under GRH. Analogous bounds for function fields of curves defined over a finite field are also shown.

We also study the distribution of values of higher derivatives of $\mathcal{L}(s, \chi) = L'(s, \chi)/L(s, \chi)$ at $s = 1$ where χ ranges over all non-trivial Dirichlet characters with a given large prime conductor m . In particular, we compute moments, i.e. the average of $P^{(a,b)}(\mathcal{L}^{(n)}(1, \chi))$, where $P^{(a,b)}(z) = z^a \bar{z}^b$ and study their asymptotic behaviour as $m \rightarrow \infty$. We then construct a density function $M_\sigma(z)$, for $\sigma = \text{Re}(s)$ and show that for $\text{Re}(s) > 1$

$$\text{Avg}_\chi \Phi(\mathcal{L}'(s, \chi)) = \int_{\mathbb{C}} M_\sigma(z) \Phi(z) |dz|$$

holds for any continuous function Φ on \mathbb{C} .

To my parents.

ACKNOWLEDGEMENTS

With heartfelt appreciation, I commence this thesis by expressing profound gratitude to my Ma. Beyond her unwavering love, I am indebted to her for being my very first math teacher, introducing me to the wonders of this subject at an early age. I truly think that her invaluable guidance has set the foundation for my academic journey. I also thank my father, my brother, and my grandmother for their continuous love and support.

I express my sincerest gratitude to my mentor and advisor Prof. V. Kumar Murty. Your guidance, support and constant motivation throughout my PhD, goes well beyond the realm of mathematics. Our numerous philosophical discussions have played a pivotal role in my personal growth, my ideas and understanding of the world and of myself.

I thank Prof. Jacob Tsimmerman and Prof. Florian Herzig for being in my supervisory committee. Their valuable comments throughout the past few years have helped shape this thesis. I also thank Prof. Arul Shankar and Prof. Steve Kudla for their exciting comments and discussions on this thesis. I am grateful to Prof. David Rohrlich who dedicated his time to thoroughly review my thesis and provided us an extensive report. Your insightful questions, comments, and suggestions are deeply appreciated. I also thank you for agreeing to be the external examiner during my defense.

I am deeply thankful to Gaurav, Hubert, Shuyang, Debanjana, Ren, Anup, Waleed, Payman, Debaditya, Vandita and in case I missed anyone, all past and present members of the GANITA Lab. You folks are amazing and over the years I have learnt a lot of math from all of you. Thank you for the discussions, being an amazing audience during my talks, asking questions and overall creating such a wonderful environment.

I also thank Stefan, Ahmee, Étienne, Sina, Eric, and Hollis for their friendships and making these years in Toronto very enjoyable. My special thanks goes to Pritam, Nilay and Shamik for their company during my undergrad days and their continued friendship.

I thank Prof. Joe Repka and Dr. Jason Siefken for their valuable feedback on my teaching. I would also like to extend my gratitude to Dr. Jonathan Korman, Prof. Marco Gualtieri, Prof. Kumar Murty and Prof. Jacob Tsimerman as I have gained a wealth of knowledge and expertise in teaching during my time as their TA.

I thank all the staff members of the department of mathematics, University of Toronto, especially Jemima, Sonja, Ashley, Patrina, Andrea and Marco. I have lost count on how many times I have dropped by at Jemima's office with some administrative issue and she happily solved them all. I also thank the staff members of The Fields Institute especially Alice and Miriam.

I am grateful for the financial support I received from the Faculty of Arts and Science, University of Toronto through their Program-level Fellowship, as well as their Alumni and Friends Graduate scholarship. I also greatly appreciate the tuition fee waivers and conference, travel and other grants, that I have received from the department over the past years. They have made my life easier and played a significant role in my graduate life experiences.

I am truly thankful to Swami Kripamayananda of Vedanta Society of Toronto for his spiritual guidance and his wonderful classes on Vivekananda's teachings. Lastly, I would like to express my deepest gratitude to Sadhguru for imparting the meditation process that changed my life! Meeting you has been the most significant event during my time in Toronto.

CONTENTS

1	INTRODUCTION	1
1.1	Organization of Chapters	1
1.2	Some background and motivation	2
1.3	Statement of main results	6
2	HIGHER EULER-KRONECKER CONSTANTS : NUMBER FIELD CASE	11
2.1	Preliminaries	11
2.2	Some background : Ihara's work on $\gamma_{K,0}$	12
2.3	Setting the stage for $\gamma_{K,1}$	14
2.4	Bounds for $\gamma_{K,1}$ under GRH	22
2.5	An unconditional arithmetic formula for $\gamma_{K,1}$	27
2.6	Generalizations to higher constants $\gamma_{K,m}$	30
2.7	Bounds for $\gamma_{K,m}$ under GRH	39
2.8	An unconditional arithmetic formula for $\gamma_{K,m}$	43
3	BOUNDS FOR THE FUNCTION FIELD CASE	45
3.1	Preliminaries	45
3.2	Bounds for $\gamma_{K,1}$	47
3.3	General Case : Bounds for $\gamma_{K,n}$	50
4	MOMENTS OF HIGHER DERIVATIVES OF $\mathcal{L}(s, \chi)$ AT $s = 1$	54
4.1	Preliminaries	54
4.2	An "exact formula" for $\mathcal{L}'(1, \chi)$	56
4.3	Generalization to higher derivatives	61
4.4	Moments : a brief history	63
4.5	Moments of $\mathcal{L}'(1, \chi)$ (conditional : under GRH)	64
4.6	Moments of $\mathcal{L}'(1, \chi)$ (unconditional)	72
4.7	Moments of higher derivatives $\mathcal{L}^{(n)}(1, \chi)$	81
5	DISTRIBUTION	86
5.1	Preliminaries	86
5.2	Distribution Functions : Some Background	88
5.3	Construction of $M_{\sigma,p}$ functions	90

5.4	A note on higher derivatives	95
5.5	Possible extension of our result to $\frac{1}{2} < \sigma \leq 1$	95
	BIBLIOGRAPHY	97

INTRODUCTION

Since the advent of analytic number theory, the study of zeros of zeta and L -functions have engaged number theorists, perhaps more than any other theme. The famous Riemann hypothesis has been an open problem for more than 160 years now. A new approach to this problem was presented by Xian-Jin Li [Li97] in 1997, who showed that positivity of a sequence of coefficients coming from the Laurent series expansion, about $s = 1$, of the logarithmic derivative of these zeta functions is equivalent to the Riemann hypothesis. Later, Brown in [Bro05] proved an effective version of Li's criterion relating positivity of the first finitely many terms in the sequence, to zero-free regions. This thesis is concerned with studying these higher coefficients.

1.1 ORGANIZATION OF CHAPTERS

In this introductory Chapter 1, we give a brief review of some well known results that will be useful for our later journey, as well as present the key motivation that led to this study. We then present a summary of our main results of this thesis.

Chapter 2 is about studying the higher *Euler-Kronecker constants* (to be defined in the next section) of a number field. In particular, after presenting some preliminary facts and Ihara's work on the constant term in sections 2.1 and 2.2 we focus on the first Euler-Kronecker constant in section 2.3. We then derive upper bounds (under GRH) in section 2.4. In 2.5 we derive, unconditionally, an arithmetic formula satisfied by this constant. In the subsequent sections 2.6 etc., we generalize these results, deduced in the previous sections, for higher constants.

Chapter 3 is a similar study in the case of a function field of a curve defined over a finite field. We prove analogous bounds.

In Chapter 4 we focus on Dirichlet L -functions. Again, section 4.1 - 4.3 is focused on deriving similar arithmetic formulas for these coefficients. The key new results in this chapter are on moments. After giving some historical background in section 4.4, we compute moments of $\mathcal{L}'(1, \chi)$ in section 4.5 under GRH, where $\mathcal{L}(s, \chi) = L'(s, \chi)/L(s, \chi)$. We then use zero sum estimates to prove an unconditional version of our result in section 4.6. Finally, we generalize these results to moments of higher derivatives in section 4.7.

Chapter 5 is on distribution of values of these higher derivatives of the logarithmic derivative of Dirichlet L -functions near $s = 1$. The main result is in section 5.3, and it is about showing the existence of a distribution function for the first derivative, for $\text{Re}(s) > 1$. In section 5.4 we briefly discuss potential generalization to higher derivatives. Finally in the concluding section 5.5 we discuss future work and issues on extension of our result to parts of the critical strip : $\frac{1}{2} < \text{Re}(s) \leq 1$.

1.2 SOME BACKGROUND AND MOTIVATION

Let K be an algebraic number field of finite degree n_K over \mathbb{Q} . The Dedekind zeta function of K is defined as

$$\zeta_K(s) = \sum_{\mathfrak{a}} (N\mathfrak{a})^{-s}$$

for $\text{Re}(s) > 1$, where the sum is taken over all integral ideals \mathfrak{a} of \mathcal{O}_K , the ring of integers of K . It also satisfies the Euler product formula

$$\zeta_K(s) = \prod_{\mathfrak{p}} \left(1 - \frac{1}{N\mathfrak{p}^s}\right)^{-1}$$

Hecke showed that $(s-1)\zeta_K(s)$ extends to an entire function. There is a simple pole of $\zeta_K(s)$ at $s=1$ and the residue satisfies the famous *class number formula* :

$$\lim_{s \rightarrow 1} (s-1)\zeta_K(s) = \frac{2^{r_1} (2\pi)^{r_2} h R}{\omega \sqrt{|d_K|}}$$

where r_1 denotes the number of real embeddings of K , $2r_2$ is that of complex embeddings, h is the class number, R is the regulator, ω is the number of roots of unity, and d_K is the discriminant of K .

For the logarithmic derivative one can write :

$$-\frac{\zeta'_K(s)}{\zeta_K(s)} = \sum_{\mathfrak{a}} \frac{\Lambda(\mathfrak{a})}{N\mathfrak{a}^s} \quad (1.1)$$

$\Lambda(\cdot)$ being the number field analogue of the von Mangoldt function given by :

$$\Lambda(\mathfrak{a}) = \begin{cases} \log N\mathfrak{p} & \text{if } \mathfrak{a} = \mathfrak{p}^k \text{ for some prime ideal } \mathfrak{p} \\ 0 & \text{otherwise.} \end{cases}$$

For the sake of completeness, we also recall that by applying the Tauberian theorem to the above (1.1), one can deduce the number field analogue of the prime number theorem, namely the prime ideal theorem :

Theorem 1.2.1. Let $\pi_K(x)$ be the number of prime ideals of \mathcal{O}_K with norm less than or equal to x . Then

$$\pi_K(x) \sim \frac{x}{\log x} \quad \text{as } x \rightarrow \infty$$

For details on the above discussion, one may refer to any standard textbook on analytic number theory, for example [Davoo], [CF76] or [MM97].

The **generalized Riemann hypothesis** (GRH) states that all non-trivial zeros (i.e. those in the critical strip) of the Dedekind zeta function is on the $s = \frac{1}{2}$ line.

Consider the (analytic) completed zeta function :

$$\tilde{\zeta}_K(s) = s(s-1)2^{r_2} \left(\frac{\sqrt{|d_K|}}{2^{r_2} \pi^{n/2}} \right)^s \Gamma\left(\frac{s}{2}\right)^{r_1} \Gamma(s)^{r_2} \zeta_K(s)$$

where $[K : \mathbb{Q}] = n_K$. In [Li97], Li introduced the following sequence of numbers $\{\lambda_n\}$, now known as Li's coefficients :

$$\lambda_n = \frac{1}{(n-1)!} \frac{d^n}{ds^n} \left(s^{n-1} \log \zeta_K(s) \right) \Big|_{s=1} \quad \text{for } n \geq 1 \quad (1.2)$$

and showed the following theorem

Theorem 1.2.2. (Li's Criterion) The general Riemann hypothesis for $\zeta_K(s)$ holds iff λ_n is non-negative for all $n \geq 1$.

Later Bombieri and Lagarias also gave an alternative proof in [BL99]. Andrew Droll, in his PhD thesis formulated a much more generalized Li's Criterion for generalized quasi-Riemann hypothesis for functions in an extension of the Selberg class.

Brown in [Bro05] proved an effective version of Li's theorem, showing positivity of the first few λ_i 's, give zero-free regions of a certain shape around $s = 1$. In particular he showed, just $\lambda_2 \geq 0$ implies non-existence of the exceptional Siegel zeros. We recall, a well-known result of Stark says that for $0 < c < \frac{1}{4}$, $\zeta_K(s)$ has at most one zero in the region

$$1 - \frac{c}{\log d_K} \leq \sigma \leq 1, \quad |t| \leq \frac{c}{\log d_K}$$

where $s = \sigma + it$. This zero, if it exists, is necessarily real and simple. We call this an exceptional Siegel zero.

Note that, if we write the Laurent series about $s = 1$ of the logarithmic derivative of $\zeta_K(s)$, then λ_2 involves the constant term and the first coefficient (that is the coefficient of $(s-1)$) together with some terms coming from the Γ -factors. This was our primary motivation to closely study this first coefficient. We later found that many of our results easily generalized to higher coefficients. A path was already led out by Ihara et. al. who, in a series of papers, systematically studied the constant term, which he called the *Euler-Kronecker constant*. We define

Definition 1.2.3. Let the Laurent series of the logarithmic derivative of $\zeta_K(s)$ about $s = 1$ be given by

$$\frac{\zeta'_K(s)}{\zeta_K(s)} = \frac{-1}{s-1} + \gamma_{K,0} + \sum_{m=1}^{\infty} \gamma_{K,n}(s-1)^m \quad (1.3)$$

$\gamma_{K,m}$ will be called the *m-th Euler-Kronecker constant*.

Remark 1.2.4. It is worth pointing out that this thesis does not deal with the subtleties of sign of these coefficients as Li's criterion demands. Instead we seek to motivate the readers to study them. These coefficients are coming from very local information, only at $s = 1$. And somehow they are able to capture what is happening at $s = \frac{1}{2}$, giving us information about all zeros! We think, in future one might be able to deduce zero-free regions and other interesting results from bounds on them and not just signs. This thesis therefore seeks to present a preliminary study of these constants.

The first thing we showed were certain arithmetic formulas for $\gamma_{K,m}$. The author was later made aware that similar formulas for $\zeta(s)$ and $\zeta_K(s)$ exists in the literature and so, we end this section with a few of those.

Suppose we write

$$\zeta(s) = \frac{1}{s-1} + \gamma + \sum_{n=1}^{\infty} \mathfrak{s}_n (s-1)^n$$

In 1885, T. J. Stieltjes [HS05] showed that

$$\mathfrak{s}_n = \frac{(-1)^n}{n!} \lim_{x \rightarrow \infty} \left(\sum_{m=1}^x \frac{(\log m)^n}{m} - \frac{(\log x)^{n+1}}{n+1} \right)$$

These \mathfrak{s}_n are called the Stieltjes constants, the generalized Euler constants or sometimes the Euler-Stieltjes constants. For the Dedekind zeta function, let us write

$$\zeta_K(s) = \sum_{n=-1}^{\infty} \mathfrak{s}_{K,n} (s-1)^n$$

The author found a similar formula in a much recent paper and does not know if similar formula has been written down in the past. The following is due to Eddin, see Theorem 2 of [Edd18].

$$\mathfrak{s}_{K,n} = \frac{(-1)^n}{n!} \lim_{x \rightarrow \infty} \left(\sum_{N\mathfrak{a} \leq x} \frac{(\log N\mathfrak{a})^n}{N\mathfrak{a}} - \mathfrak{s}_{K,-1} \frac{(\log x)^{n+1}}{n+1} \right) \quad \text{for } n \geq 1$$

and

$$\mathfrak{s}_{K,0} = \lim_{x \rightarrow \infty} \left(\sum_{N\mathfrak{a} \leq x} \frac{1}{N\mathfrak{a}} - \mathfrak{s}_{K,-1} \log x \right) + \mathfrak{s}_{K,-1}$$

The following formula for the logarithmic derivative of the Riemann zeta function is also known. Let

$$\frac{\zeta'(s)}{\zeta(s)} = \frac{-1}{(s-1)} + \sum_{n=0}^{\infty} \gamma_n (s-1)^n$$

then

$$\gamma_n = \frac{(-1)^{n-1}}{n!} \lim_{x \rightarrow \infty} \left(\sum_{m < x} \frac{\Lambda(m)(\log m)^n}{m} - \frac{(\log x)^{n+1}}{n+1} \right)$$

For a proof see [Tit58].

The author is not aware of existence of a similar formula for Dedekind zeta functions. The formulas we deduced are similar but a bit more involved.

1.3 STATEMENT OF MAIN RESULTS

Our first result is the following formula :

Theorem 1.3.1. (Unconditionally)

$$\gamma_{K,1} = \lim_{x \rightarrow \infty} \left[\Psi_K(x) - 1 - \frac{1+x}{1-x} \log x - \frac{1}{2} (\log x)^2 \right]$$

$$\text{where, } \Psi_K(x) = \frac{1}{x-1} \sum_{k, N(P)^k \leq x} \left(\frac{x}{N(P)^k} - 1 \right) k (\log N(P))^2 \quad \text{for } x > 1$$

Theorem 1.3.2. Under GRH, for $|d_K| > 8$ and writing $\alpha_K = \log \sqrt{|d_K|}$, we have

$$|\gamma_{K,1}| \leq 2(\log \alpha_K)(2 \log \alpha_K - \gamma_{K,0}) + 18 \log \alpha_K + O\left(\frac{(\log \alpha_K)^2}{\alpha_K}\right)$$

Corollary 1.3.3. When $\gamma_{K,0} \geq 0$, we get, (under GRH and for $|d_K| > 8$)

$$\gamma_{K,1} \ll \left(\log \log \sqrt{|d_K|} \right)^2$$

Similarly for the general case we write :

$$\Psi_K(m, x) = \frac{1}{x-1} \sum_{k, N(P)^k \leq x} \left(\frac{x}{N(P)^k} - 1 \right) k^m (\log N(P))^{m+1} \quad \text{for } x > 1$$

Theorem 1.3.4. (Unconditionally)

$$\gamma_{K,m} + (-1)^m = \lim_{x \rightarrow \infty} \frac{1}{m!} \left[(-1)^{m+1} \Psi_K(m, x) + \frac{f(m, x)}{(x-1)} \right] \quad (1.4)$$

where $f(m, x)$ is recursively defined as :

$$f(m, x) = \frac{(-1)^m}{m+1} (x-1)(\log x)^{m+1} + (-1)^{m+1} (x+1)(\log x)^m + m(m-1)f(m-2, x)$$

$$f(1, x) = (1-x) \left[2 + \frac{1+x}{1-x} \log x + \frac{1}{2} (\log x)^2 \right]$$

$$f(0, x) = (x-1) \log(x)$$

Theorem 1.3.5. Under GRH, for $|d_K| > 8$, and $m \geq 1$ we have

$$\gamma_{K,m} \ll \frac{2^m}{m!} (\log(2^m(m!)^2) + \log \alpha_K)^m (2 \log(2^m(m!)^2) + 2 \log \alpha_K - \gamma_{K,0} + 1)$$

where as before, $\alpha_K = \log \sqrt{|d_K|}$.

As a corollary we have :

Corollary 1.3.6. When $\gamma_{K,0} \geq 0$ we get, (under GRH and for $|d_K| > 8$)

$$\gamma_{K,m} \ll \frac{2^{m+1}}{m!} (\log(2^m(m!)^2) + \log \alpha_K)^{m+1}$$

In particular, for $m \ll \frac{\log d_K}{\log \log d_K}$, we have

$$\gamma_{K,m} \ll \frac{2^{m+1}}{m!} (\log \log d_K)^{m+1}$$

We also prove analogous bounds in the function field case (unconditional, GRH being known). For example if K is the function field of a curve X of genus g , over \mathbb{F}_q , q being a prime power then :

Theorem 1.3.7. For $g > 2$ or, $g = 2$ and $q > 2$, and writing $\alpha_K = (g - 1) \log q$, we have

$$\gamma_{K,n} + (-1)^n \ll \frac{2}{n!} (\log(n!2^{n+1}\alpha_K))^n (2 \log(\alpha_K) - \gamma_{K,0} + \log q + 1 + n!2^n)$$

Remark 1.3.8. All implicit constants in the above results are absolute.

Remark 1.3.9. Just for consistency and clarity of notation, let us mention that, by $a \ll b$ we mean that there exists a positive constant c such that $|a| \leq cb$. Sometimes we also might have used the big- O notation. They mean the same thing, i.e. $a = O(b) \iff a \ll b$.

We then turn our attention to distribution of values of these higher coefficients. For this we consider the following setting : let K be a number field and χ be a primitive Dirichlet character on K . Let $L(s, \chi)$ be the L -function associated to it. For notational brevity we'll write

$$\mathcal{L}(s, \chi) := \frac{L'(s, \chi)}{L(s, \chi)}$$

Initially we proved similar formulae and bounds like that of $\gamma_{K,n}$, namely,

$$\mathcal{L}^{(n)}(1, \chi) = \lim_{x \rightarrow \infty} (-1)^{n+1} \Psi_K(\chi, n, x)$$

where

$$\Psi_K(\chi, n, x) = \frac{1}{x-1} \sum_{k, N(P)^k < x} k^n \left(\frac{x}{N(P)^k} - 1 \right) \chi(P)^k (\log N(P))^{n+1}$$

But our main goal was to compute moments for the higher derivatives, motivated by the work of Ihara, Murty and Shimura, who in [IMS09], computed moments of $\mathcal{L}(1, \chi)$.

For this section we work with $K = \mathbb{Q}$. Let m be a prime and X_m denote the set of all non-principal multiplicative characters $\chi : (\mathbb{Z}/m\mathbb{Z})^\times \rightarrow \mathbb{C}^\times$ and $L(s, \chi)$ denote the corresponding Dirichlet L -function. For any pair of

non-negative integers (a, b) let $P^{(a,b)}(z) = z^a \bar{z}^b$. We showed :

Theorem 1.3.10. For any $\epsilon > 0$ we have, unconditionally,

$$\frac{1}{|X_m|} \sum_{\chi \in X_m} P^{(a,b)}(\mathcal{L}^{(r)}(1, \chi)) = (-1)^{(r+1)d} \mu^{(a,b)}(r) + O\left(m^{\epsilon-1}\right)$$

The implicit constant depends on a, b only. Under GRH, the error term is

$$O\left(\frac{(\log m)^{(r+1)d+2}}{m}\right)$$

with $d = a + b$. In particular, letting $m \rightarrow \infty$ we get

$$\lim_{m \rightarrow \infty} \frac{1}{|X_m|} \sum_{\chi \in X_m} P^{(a,b)}(\mathcal{L}^{(r)}(1, \chi)) = (-1)^{(r+1)d} \mu^{(a,b)}(r)$$

Here $\mu^{(a,b)}(r)$ is the following explicitly computable constant :

$$\mu^{(a,b)}(r) = \sum_{j=1}^{\infty} \frac{\ell^r \Lambda_a(j) \ell^r \Lambda_b(j)}{j^2}$$

where for $k > 0, r \geq 0$

$$\ell^r \Lambda_k(n) = \sum_{n_1 n_2 \cdots n_k = n} \left(\prod_{i=1}^k \Lambda(n_i) (\log n_i)^r \right)$$

whereas, for $k = 0, \ell^r \Lambda_0(n) = 1$ for $n = 1$ and 0 for $n > 1$.

Note : $\ell^r \Lambda_a(\cdot)$ is just notation. We are not actually multiplying by some ℓ^r or anything. The logarithm appears with exponent r in the above formula together with $\Lambda(\cdot)$, this is just a way of book keeping.

We then focus on the distribution of values of these higher derivatives of the logarithmic derivative of Dedekind zeta functions, in particular we show the existence of a density function for $\text{Re}(s) > 1$.

Let K be either \mathbb{Q} or an imaginary quadratic number field. Let χ run over all Dirichlet characters on K whose conductor (the non-archimedean part) is a prime divisor, such that $\chi(\wp_\infty) = 1$. We define the average of

a complex valued function $\phi(\chi)$, over a family of χ as defined above, as follows :

$$\text{Avg}_\chi \phi(\chi) = \lim_{m \rightarrow \infty} \text{Avg}_{N(\mathfrak{f}) \leq m} \phi(\chi)$$

where

$$\text{Avg}_{N(\mathfrak{f}) \leq m} \phi(\chi) = \frac{\sum_{N(\mathfrak{f}) \leq m} \left(\sum_{\chi = \mathfrak{f}} \phi(\chi) \right) / \sum_{\chi = \mathfrak{f}} 1}{\sum_{N(\mathfrak{f}) \leq m} 1}$$

Then the main result in this chapter states that,

Theorem 1.3.11. For any $s \in \mathbb{C}$ with $\sigma = \text{Re}(s) > 1$ there exists a function $M_\sigma : \mathbb{C} \rightarrow \mathbb{R}$ satisfying, $M_\sigma(w) \geq 0$, and $\int_{\mathbb{C}} M_\sigma(w) |dw| = 1$, such that

$$\text{Avg}_\chi \Phi(\mathcal{L}'(\chi, s)) = \int_{\mathbb{C}} M_\sigma(w) \Phi(w) |dw| \quad (1.5)$$

holds for any continuous function Φ of \mathbb{C} .

Note that M_σ is constructed as the limit of $M_{\sigma, P}$ functions, where P is a finite set of non-archimedean primes. The Fourier dual of $M_\sigma(z)$ given by

$$\tilde{M}_\sigma(z) = \int_{\mathbb{C}} M_\sigma(w) \psi_z(w) |dw|$$

where $\psi_z : \mathbb{C} \rightarrow \mathbb{C}^1$ is the additive character $\psi_z(w) = \exp(i \cdot \text{Re}(\bar{z}w))$, satisfies the following :

$$\tilde{M}_\sigma(z) = \text{Avg}_\chi \psi_z(\mathcal{L}'(\chi, s))$$

HIGHER EULER-KRONECKER CONSTANTS : NUMBER FIELD CASE

2.1 PRELIMINARIES

Let K be an algebraic number field. The Dedekind zeta function $\zeta_K(s)$ has a simple pole at $s = 1$ and $(s - 1)\zeta_K(s)$ extends to an entire function in the complex plane. Therefore one can write a Laurent series of $\zeta_K(s)$ about $s = 1$ as follows :

$$\zeta_K(s) = \frac{c_{-1}}{s-1} + c_0 + c_1(s-1) + c_2(s-1)^2 + \dots \quad (c_{-1} \neq 0) \quad (2.1)$$

In [Ih96], Ihara studied the constant $\gamma_K = c_0/c_{-1}$ attached to K and called it the *Euler-Kronecker constant*. Note that the logarithmic derivative of $\zeta_K(s)$ also has a simple pole at $s = 1$, the key difference being, the residue is then just -1 . We can write down a Laurent series for $\zeta'_K(s)/\zeta_K(s)$ about $s = 1$, turns out the constant of this series is γ_K . Our work in this chapter is to analyze the coefficients of higher powers of $(s - 1)$. We will refer to these as general or higher Euler-Kronecker constants.

Definition 2.1.1. Let the Laurent series of the logarithmic derivative of $\zeta_K(s)$ about $s = 1$ be given by

$$\frac{\zeta'_K(s)}{\zeta_K(s)} = \frac{-1}{s-1} + \gamma_{K,0} + \sum_{m=1}^{\infty} \gamma_{K,m}(s-1)^m \quad (2.2)$$

$\gamma_{K,m}$ will be called the *m-th Euler-Kronecker constant*.

For $K = \mathbb{Q}$, the constant term is the famous Euler-Mascheroni constant

$$\gamma_{\mathbb{Q},0} = \gamma = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n} - \log n \right) = 0.57721566 \dots$$

We now describe some results obtained by Ihara on $\gamma_{K,0}$.

2.2 SOME BACKGROUND : IHARA'S WORK ON $\gamma_{K,0}$

We first note that :

$$\gamma_{K,0} = \lim_{s \rightarrow 1} \left(\frac{\zeta'_K(s)}{\zeta_K(s)} + \frac{1}{s-1} \right)$$

On the other hand using a lemma of Stark, (Lemma 3 in [Sta74]) we get

$$-\frac{\zeta'_K(s)}{\zeta_K(s)} = \frac{1}{s} + \frac{1}{s-1} - \sum \frac{1}{s-\rho} + \alpha_K + \beta_K + \tilde{\Gamma}_K(s) \quad (2.3)$$

where the sum runs over all non-trivial zeros ρ of $\zeta_K(s)$, counted with multiplicities. Note that this result is just a consequence of Hadamard factorization. Also, in the above,

$$\alpha_K = \frac{1}{2} \log |d_K|, \quad d_K \text{ being the absolute discriminant of } K ;$$

$$\beta_K = - \left\{ \frac{r_1}{2} (\gamma + \log 4\pi) + r_2 (\gamma + \log 2\pi) \right\}$$

$$\tilde{\Gamma}_K(s) = \frac{r_1}{2} \left(g\left(\frac{s}{2}\right) - g\left(\frac{1}{2}\right) \right) + r_2 (g(s) - g(1)) \quad \text{where } g(s) = \frac{\Gamma'(s)}{\Gamma(s)}$$

where K has r_1 real conjugate fields and $2r_2$ complex conjugate fields, and $\gamma = \gamma_{\mathbb{Q},0}$ as before. Thus taking the $\frac{1}{s-1}$ to other side and letting $\lim s \rightarrow 1$ we get,

$$\begin{aligned} -\gamma_{K,0} &= 1 - \sum_{\rho} \frac{1}{1-\rho} + \alpha_K + \beta_K \\ &\Rightarrow \frac{1}{2} \sum_{\rho} \frac{1}{\rho(1-\rho)} = \gamma_{K,0} + \alpha_K + \beta_K + 1 \end{aligned} \quad (2.4)$$

Notice that $\tilde{\Gamma}_K(1) = 0$. Equation (2.4) will be used in a later section while finding upper bounds of certain terms.

Ihara proved the following bounds for $\gamma_{K,0}$ in [Iha06], e.g. see Theorem 1 and Proposition 3. (Although the lower bound mentioned below is trivial, as the left hand side of (2.4) above is positive.)

Theorem 2.2.1. (Ihara) For $n_K = [K : \mathbf{Q}] > 2$ or, $n_K = 2$ and $|d_K| > 8$, we have (the constants c_1, c_2 below are absolute)

$$\begin{aligned} \gamma_{K,0} &\leq c_1 \log \log \sqrt{|d_K|} \quad (\text{under GRH}) \\ &\geq -c_2 \log \sqrt{|d_K|} \quad (\text{unconditionally}) \end{aligned}$$

To demonstrate that the general (negative) lower bound cannot be so close to 0 as the upper we note that

$$-0.26049\dots \leq \liminf \frac{\gamma_{K,0}}{\log \sqrt{|d_K|}} \leq -0.17849\dots$$

The left side inequality is under GRH, the right side is unconditional, both due to Tsfasman e.g. see [Tsf06].

We quote a few other results from Ihara's paper [Iha06], as we will either use them in subsequent sections or prove analogous versions for higher Euler-Kronecker constants. Consider the prime counting function

$$\Phi_K(x) = \frac{1}{x-1} \sum_{k, N(P)^k \leq x} \left(\frac{x}{N(P)^k} - 1 \right) \log N(P) \quad (\text{for } x > 1)$$

where P runs over non-archimedean primes of K and k over positive integers such that $N(P)^k \leq x$. For large x , this function behaves like $\log x$, in fact Ihara shows the following formula unconditionally :

$$\lim_{x \rightarrow \infty} (\log x - \Phi_K(x)) = \gamma_{K,0} + 1 \tag{2.5}$$

One also has the following upper bound on $\Phi_K(x)$

$$\Phi_K(x) \leq \log x - \frac{\sqrt{x}-1}{\sqrt{x}+1}(\gamma_{K,0} + 1) + \frac{2}{\sqrt{x}+1}(\alpha_K + \beta_k) + \frac{n_K(\log x + 1)}{x-1} \tag{2.6}$$

This is a consequence of Main Lemma (see 1.5.6) and Lemma 2 of [Iha06].

In the subsequent sections, we will first investigate the next coefficient $\gamma_{K,1}$ and will then try to see whether the methods used can be generalized.

2.3 SETTING THE STAGE FOR $\gamma_{K,1}$

Recall, we wrote

$$\frac{\zeta'_K(s)}{\zeta_K(s)} + \frac{1}{s-1} = \gamma_{K,0} + \sum_{m=1}^{\infty} \gamma_{K,m}(s-1)^m \quad (2.7)$$

Taking derivative and letting $\lim s \rightarrow 1$ we get,

$$\lim_{s \rightarrow 1} \left[\frac{d}{ds} \frac{\zeta'_K(s)}{\zeta_K(s)} - \frac{1}{(s-1)^2} \right] = \gamma_{K,1} \quad (2.8)$$

From the Euler product, one has

$$-\frac{\zeta'_K(s)}{\zeta_K(s)} = \sum_{P, k \geq 1} \frac{\log N(P)}{N(P)^{ks}}$$

For brevity of notation, we will denote the left hand side by $Z_K(s)$, i.e.

$$Z_K(s) = -\frac{\zeta'_K(s)}{\zeta_K(s)}$$

Thus taking derivative, one has

$$Z'_K(s) = \sum_{P, k \geq 1} \frac{-k(\log N(P))^2}{N(P)^{ks}} \quad (2.9)$$

On the other hand, differentiating the expression obtained from the Hadamard product as in (2.3) we get,

$$Z'_K(s) = -\frac{1}{s^2} - \frac{1}{(s-1)^2} + \sum \frac{1}{(s-\rho)^2} + \tilde{\Gamma}'_K(s) \quad (2.10)$$

with $\tilde{\Gamma}'_K(s) = \frac{r_1}{4} g'(\frac{s}{2}) + r_2 g'(s)$, where, as before $g(s) = \frac{\Gamma'}{\Gamma}(s)$.

Taking the $\frac{1}{(s-1)^2}$ on the other side and letting $\lim s \rightarrow 1$, the left hand side becomes $-\gamma_{K,1}$. Thus we have

$$\boxed{\gamma_{K,1} = 1 - \sum \frac{1}{(1-\rho)^2} - \tilde{\Gamma}'_K(1)} \quad (2.11)$$

We wish to find a similar arithmetic formula as in (2.5) and bounds for $\gamma_{K,1}$. To do so, we consider the integral

$$\Psi_K^{(\mu)}(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^{s-\mu}}{s-\mu} Z'_K(s) ds \quad \text{for } c \gg 0$$

for $\mu = 0$ and 1 and evaluate the expression $x\Psi_K^{(1)}(x) - \Psi_K^{(0)}(x)$ in two different ways using equation (2.9) and equation (2.10). Note that by $c \gg 0$, we just mean that we are considering the integral on a line $s = c$, far to the right of 1 .

The following classical formulas will be of help.

$$\frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{y^s}{s} ds = \begin{cases} 0 & 0 < y < 1 \\ \frac{1}{2} & y = 1 \\ 1 & y > 1 \end{cases} \quad (2.12)$$

And for $n \geq 1$,

$$\frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{y^s}{s^{n+1}} ds = \begin{cases} 0 & 0 < y \leq 1 \\ \frac{1}{n!} (\log y)^n & y > 1 \end{cases} \quad (2.13)$$

Using the expression for $Z'_K(s)$ from (2.9) we get :

$$\begin{aligned} x\Psi_K^{(1)}(x) - \Psi_K^{(0)}(x) &= x \cdot \sum_{P,k \geq 1} \frac{-k(\log N(P))^2}{N(P)^k} \left[\frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{1}{s-1} \left(\frac{x}{N(P)^k} \right)^{s-1} ds \right] \\ &\quad - \sum_{P,k \geq 1} -k(\log N(P))^2 \left[\frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{1}{s} \left(\frac{x}{N(P)^k} \right)^s ds \right] \\ &= x \cdot \sum_{k, N(P)^k < x} \frac{-k(\log N(P))^2}{N(P)^k} + \sum_{k, N(P)^k = x} \frac{-k(\log N(P))^2}{2} \\ &\quad - \sum_{k, N(P)^k < x} -k(\log N(P))^2 - \sum_{k, N(P)^k = x} \frac{-k(\log N(P))^2}{2} \\ &= \sum_{k, N(P)^k \leq x} \left(\frac{x}{N(P)^k} - 1 \right) (-k(\log N(P))^2) \end{aligned}$$

Looking at the above computation, we define :

$$\Psi_K(x) = \frac{1}{x-1} \sum_{k, N(P)^k \leq x} \left(\frac{x}{N(P)^k} - 1 \right) k(\log N(P))^2 \quad \text{for } x > 1$$
(2.14)

Remark 2.3.1. The reason for dividing by $(x-1)$ will become apparent while computing $x\Psi_K^{(1)}(x) - \Psi_K^{(0)}(x)$ the other way. For now we have,

$$x\Psi_K^{(1)}(x) - \Psi_K^{(0)}(x) = -(x-1)\Psi_K(x) \quad (2.15)$$

Now using the expression for $Z'_K(s)$ from (2.10) we get :

$$\begin{aligned} x\Psi_K^{(1)}(x) - \Psi_K^{(0)}(x) &= \frac{x}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^{s-1}}{s-1} \left[-\frac{1}{s^2} - \frac{1}{(s-1)^2} + \sum \frac{1}{(s-\rho)^2} + \tilde{\Gamma}'_K(s) \right] ds \\ &\quad - \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^s}{s} \left[-\frac{1}{s^2} - \frac{1}{(s-1)^2} + \sum \frac{1}{(s-\rho)^2} + \tilde{\Gamma}'_K(s) \right] ds \end{aligned}$$

Let us first look at the contribution from the term : $-\frac{1}{s^2} - \frac{1}{(s-1)^2}$

$$\begin{aligned} &\int x^s \left[-\frac{1}{s^2(s-1)} - \frac{1}{(s-1)^3} + \frac{1}{s^3} + \frac{1}{s(s-1)^2} \right] ds \\ &= \int x^s \left[\frac{1}{s^2} + \frac{1}{s} - \frac{1}{(s-1)} - \frac{1}{(s-1)^3} + \frac{1}{s^3} + \frac{1}{(s-1)^2} - \frac{1}{s-1} + \frac{1}{s} \right] ds \\ &= 2 \int \frac{x^s}{s} ds - 2x \int \frac{x^{s-1}}{s-1} ds + \int x^s \left[\frac{1}{s^2} + \frac{1}{(s-1)^2} \right] ds + \int x^s \left[\frac{1}{s^3} - \frac{1}{(s-1)^3} \right] ds \end{aligned}$$

(Thus using the classical formulas as in (2.12) and (2.13) we have)

$$\begin{aligned} &= 2 - 2x + \log x(1+x) + \frac{1}{2}(\log x)^2(1-x) \\ &= (1-x) \left[2 + \frac{1+x}{1-x} \log x + \frac{1}{2}(\log x)^2 \right] := f(x) \quad (\text{say}) \end{aligned}$$
(2.16)

Note that

$$\frac{f(x)}{(x-1)^2} \ll (\log x)^2 \tag{2.17}$$

Now we focus on the contribution from the \sum_ρ term. Recall we're trying to evaluate

$$\frac{x}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^{s-1}}{s-1} \sum \frac{1}{(s-\rho)^2} ds - \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^s}{s} \sum \frac{1}{(s-\rho)^2} ds$$

We will do some contour manipulation for this. As in Figure 2.1 (below), for large T and R (to be chosen later), take the contour $C_{R,T}$ to be the rectangle : $c - iT \rightarrow c + iT \rightarrow -R + iT \rightarrow -R - iT \rightarrow c - iT$.




Figure 2.1

We're interested only in the side $c - iT \rightarrow c + iT$. We'll show that for specific choice of R , the contribution from the other three sides of the rectangle goes to 0 as $T \rightarrow \infty$.

On the side $\{-R + iT \rightarrow -R - iT\}$, writing $\rho = \beta + i\gamma$ and $s = -R + it$ and so, $ds = idt$ we have,

$$\begin{aligned} \left| \int_{-R+iT}^{-R-iT} \frac{x^s}{s} \cdot \frac{1}{(s-\rho)^2} ds \right| &\leq \int_{-T}^T \frac{x^{-R}}{\sqrt{R^2 + t^2}((R+\beta)^2 + (t-\gamma)^2)} dt \\ &\leq \frac{x^{-R}}{R^3} \int_{-T}^T dt \leq 2x^{-R} \frac{T}{R^3} \end{aligned}$$

Similarly, for the integral on $\{c + iT \rightarrow -R + iT\}$, writing $s = \sigma + iT$

$$\begin{aligned} \left| \int_{c+iT}^{-R+iT} \frac{x^s}{s} \cdot \frac{1}{(s-\rho)^2} ds \right| &\leq \int_c^{-R} \frac{x^\sigma}{\sqrt{\sigma^2 + T^2}((\sigma-\beta)^2 + (T-\gamma)^2)} d\sigma \\ &\leq \frac{x^c}{T} \int_c^{-R} \frac{d\sigma}{(\sigma-\beta)^2} = \frac{x^c}{T} \left[\frac{-1}{\sigma-\beta} \right]_c^{-R} \\ &= \frac{x^c}{T} \left[\frac{-1}{-R-\beta} + \frac{1}{c-\beta} \right] = \frac{x^c}{T} \left[\frac{1}{R+\beta} + \frac{1}{c-\beta} \right] \\ &= \frac{x^c}{T} \frac{c+R}{(R+\beta)(c-\beta)} \ll \frac{x^c}{T} \end{aligned}$$

Note that the last inequality follows from, $0 < \beta < 1$ and we can choose $c \geq 2$, so that $c - \beta \geq 1$, whereas, $\frac{c+R}{R+\beta} \ll 1$.

Similarly, writing $s = \sigma - iT$, we have

$$\begin{aligned} \left| \int_{-R-iT}^{c-iT} \frac{x^s}{s} \cdot \frac{1}{(s-\rho)^2} ds \right| &\leq \int_{-R}^c \frac{x^\sigma}{\sqrt{\sigma^2 + T^2}((\sigma-\beta)^2 + (T+\gamma)^2)} d\sigma \\ &\leq \frac{x^c}{T} \int_c^{-R} \frac{d\sigma}{(\sigma-\beta)^2} \ll \frac{x^c}{T} \end{aligned}$$

as before. Thus by choosing $R = T$ and letting, $T \rightarrow \infty$ we see that these integrals go to zero. Therefore by residue theorem, the line integral on $s = c$ is same as the residue at the poles to the left of c .

Now let us compute these residues.

The pole at $s = 0$ has residue : $\frac{1}{\rho^2}$.

The pole at ρ (double pole) has residue : $\lim_{s \rightarrow \rho} \frac{d}{ds} \left(\frac{x^s}{s} \right) = \frac{\rho x^\rho \log x - x^\rho}{\rho^2}$
(Computations for $s = 1$ are exactly similar).

Therefore by Residue theorem, net contribution from the \sum_ρ term is given by :

$$\begin{aligned}
& x \left[\sum \frac{1}{(\rho-1)^2} + \sum \frac{(\rho-1)x^{\rho-1} \log x - x^{\rho-1}}{(\rho-1)^2} \right] - \left[\sum \frac{1}{\rho^2} + \sum \frac{\rho x^\rho \log x - x^\rho}{\rho^2} \right] \\
&= (x-1) \sum \frac{1}{(1-\rho)^2} + \sum \frac{\rho^2(\rho-1)x^\rho \log x - \rho^2 x^\rho - \rho(\rho-1)^2 x^\rho \log x + (\rho-1)^2 x^\rho}{\rho^2(\rho-1)^2} \\
&= (x-1) \sum \frac{1}{(1-\rho)^2} + \sum \frac{\rho(\rho-1)x^\rho \log x(\rho-\rho+1) - \rho^2 x^\rho + (\rho-1)^2 x^\rho}{\rho^2(\rho-1)^2} \\
&= (x-1) \left[\sum \frac{\rho(\rho-1) \log x - \rho^2 + (1-\rho)^2}{\rho^2(1-\rho)^2} \cdot \frac{x^\rho}{x-1} + \sum \frac{1}{(1-\rho)^2} \right] \\
&= (x-1) \left[r(x) + \sum \frac{1}{(1-\rho)^2} \right] \quad (\text{say})
\end{aligned} \tag{2.18}$$

So we are defining everything except the $\sum \frac{1}{(1-\rho)^2}$ term, in the previous expression as $r(x)$, i.e.

$$r(x) = \sum \frac{\rho(\rho-1) \log x - \rho^2 + (1-\rho)^2}{\rho^2(1-\rho)^2} \cdot \frac{x^\rho}{x-1} \tag{2.19}$$

For the Gamma term, we first recall

$$\tilde{\Gamma}_K(s) = \frac{r_1}{2} \left[g\left(\frac{s}{2}\right) - g\left(\frac{1}{2}\right) \right] + r_2 [g(s) - g(1)]$$

where $g(s) = \frac{\Gamma'}{\Gamma}(s)$ is the *digamma* function. Thus, taking derivative we get,

$$\tilde{\Gamma}'_K(s) = \frac{r_1}{4} g'\left(\frac{s}{2}\right) + r_2 g'(s)$$

We look at the series expansion of $\frac{\Gamma'}{\Gamma}(s)$ and its derivative.

$$\begin{aligned}
\frac{\Gamma'}{\Gamma}(s) &= g(s) = -\gamma - \sum_{k=0}^{\infty} \left(\frac{1}{s+k} - \frac{1}{1+k} \right) \\
\Rightarrow g'(s) &= \sum_{k=0}^{\infty} \frac{1}{(s+k)^2}
\end{aligned}$$

Note that the above series expansion of digamma is valid in the entire complex plane except for non-positive integers. i.e. in $\mathbb{C} \setminus \{0, -1, -2, -3, \dots\}$.

Therefore we have

$$\tilde{\Gamma}'_K(s) = \frac{r_1 + r_2}{s^2} + r_1 \left(\sum_{n=1}^{\infty} \frac{1}{(s+2n)^2} \right) + r_2 \left(\sum_{n=1}^{\infty} \frac{1}{(s+n)^2} \right)$$

We want to compute :

$$\frac{x}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^{s-1}}{s-1} \tilde{\Gamma}'_K(s) ds - \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^s}{s} \tilde{\Gamma}'_K(s) ds$$

We compute the contribution from the three constituent terms of $\tilde{\Gamma}_K(s)$, as above, individually.

$$\begin{aligned} & (r_1 + r_2) \left[\frac{x}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^{s-1}}{s^2(s-1)} ds - \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^s}{s^3} ds \right] \\ &= (r_1 + r_2) \left[\frac{x}{2\pi i} \int_{c-i\infty}^{c+i\infty} x^{s-1} \left(\frac{1}{s-1} - \frac{1}{s} - \frac{1}{s^2} \right) ds - \frac{1}{2} (\log x)^2 \right] \\ &= (r_1 + r_2) \left[x - 1 - \log x - \frac{1}{2} (\log x)^2 \right] \end{aligned}$$

Note that for the two series we can again refer to Figure 2.1 and use similar residue computations as $\frac{1}{(s-\rho)^2}$ with $\rho = -2n$ and $\rho = -n$.

For example, for $\rho = -n$, the term will be :

$$r_2(x-1) \left[\sum_{n=1}^{\infty} \frac{n(n+1) \log x - n^2 + (1+n)^2}{n^2(1+n)^2} \cdot \frac{x^{-n}}{x-1} + \sum_{n=1}^{\infty} \frac{1}{(1+n)^2} \right] - r_2$$

Thus we putting these together we get,

$$\begin{aligned}
& \frac{x}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^{s-1}}{s-1} \tilde{\Gamma}'_K(s) ds - \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^s}{s} \tilde{\Gamma}'_K(s) ds \\
&= (r_1 + r_2)(x-1) \left[1 - \frac{1}{x-1} \log x - \frac{1}{2(x-1)} (\log x)^2 \right] \\
&+ r_1(x-1) \left[\sum_{n=1}^{\infty} \frac{2n(2n+1) \log x - 4n^2 + (1+2n)^2}{4n^2(1+2n)^2} \cdot \frac{x^{-2n}}{x-1} + \sum_{n=1}^{\infty} \frac{1}{(1+2n)^2} \right] \\
&+ r_2(x-1) \left[\sum_{n=1}^{\infty} \frac{n(n+1) \log x - n^2 + (1+n)^2}{n^2(1+n)^2} \cdot \frac{x^{-n}}{x-1} + \sum_{n=1}^{\infty} \frac{1}{(1+n)^2} \right] \\
&+ r_1 \left[\sum_{n=1}^{\infty} \frac{1}{(1+2n)^2} - \sum_{n=1}^{\infty} \frac{1}{4n^2} \right] - r_2 \\
&= (x-1) [\ell(x) + \tilde{\Gamma}'_K(1)] \quad (\text{say})
\end{aligned} \tag{2.20}$$

The idea here is the same as the non-trivial zero case, we are isolating $\tilde{\Gamma}'_K(1)$ and denoting the rest of the expression by $\ell(x)$.

Note that, $\tilde{\Gamma}'_K(1) = (r_1 + r_2) + r_1 \sum_{n=1}^{\infty} \frac{1}{(1+2n)^2} + r_2 \sum_{n=1}^{\infty} \frac{1}{(1+n)^2}$.

Thus,

$$\begin{aligned}
\ell(x) &= -\frac{r_1 + r_2}{x-1} \left[\log x + \frac{1}{2} (\log x)^2 \right] + \frac{r_1}{x-1} \left[\sum_{n=1}^{\infty} \frac{1}{(1+2n)^2} - \sum_{n=1}^{\infty} \frac{1}{4n^2} \right] \\
&+ r_1 \left[\sum_{n=1}^{\infty} \frac{2n(2n+1) \log x - 4n^2 + (1+2n)^2}{4n^2(1+2n)^2} \cdot \frac{x^{-2n}}{x-1} \right] \\
&+ r_2 \left[\sum_{n=1}^{\infty} \frac{n(n+1) \log x - n^2 + (1+n)^2}{n^2(1+n)^2} \cdot \frac{x^{-n}}{x-1} \right] - \frac{r_2}{x-1} \quad (2.21)
\end{aligned}$$

As we'll soon see, although the above expressions look complicated, they can be very easily estimated.

Putting equations (2.15), (2.16), (2.18), (2.20) together,

$$\begin{aligned}
-(x-1)\Psi_K(x) &= (1-x) \left[2 + \frac{1+x}{1-x} \log x + \frac{1}{2}(\log x)^2 \right] \\
&\quad + (x-1) \left[r(x) + \sum \frac{1}{(1-\rho)^2} \right] + (x-1) [\ell(x) + \tilde{\Gamma}'_K(1)] \\
\Rightarrow \Psi_K(x) &= 2 + \frac{1+x}{1-x} \log x + \frac{1}{2}(\log x)^2 - r(x) - \sum \frac{1}{(1-\rho)^2} - \ell(x) - \tilde{\Gamma}'_K(1) \\
&= \gamma_{K,1} + 1 + \frac{1+x}{1-x} \log x + \frac{1}{2}(\log x)^2 - r(x) - \ell(x)
\end{aligned}$$

Note the last equality follows from equation (2.11).

$$\boxed{\gamma_{K,1} = \Psi_K(x) - 1 - \frac{1+x}{1-x} \log x - \frac{1}{2}(\log x)^2 + r(x) + \ell(x)} \quad (2.22)$$

We are now ready to deduce some bounds.

2.4 BOUNDS FOR $\gamma_{K,1}$ UNDER GRH

Estimates for $\Psi_K(x)$

Recall,

$$\begin{aligned}
\Psi_K(x) &= \frac{1}{x-1} \sum_{k, N(P)^k \leq x} \left(\frac{x}{N(P)^k} - 1 \right) k(\log N(P))^2 \\
&= \frac{1}{x-1} \sum_{k, N(P)^k \leq x} \left(\frac{x}{N(P)^k} - 1 \right) (\log N(P))(\log N(P))^k \\
&\leq (\log x)\Phi_K(x)
\end{aligned} \quad (2.23)$$

$$\text{Where } \Phi_K(x) = \frac{1}{x-1} \sum_{k, N(P)^k \leq x} \left(\frac{x}{N(P)^k} - 1 \right) (\log N(P)).$$

This is the counterpart of our $\Psi_K(x)$ used by Ihara in [Iha06] to compute $\gamma_{K,0}$. Also note that $\Psi_K(x) \geq 0$. As noted in the previous section, e.g. see (2.6) Ihara showed the upper bound

$$\begin{aligned}
\Phi_K(x) &\leq \log x - \frac{\sqrt{x}-1}{\sqrt{x}+1}(\gamma_{K,0}+1) + \frac{2}{\sqrt{x}+1}(\alpha_K + \beta_K) + \frac{n_K(\log x + 1)}{x-1} \\
\Rightarrow \Psi_K(x) &\leq (\log x)^2 - \left(\frac{\sqrt{x}-1}{\sqrt{x}+1}\right)(\log x)(\gamma_{K,0}+1) + \frac{2\log x}{\sqrt{x}}(\alpha_K + \beta_K) \\
&\quad + \frac{n_K(\log x)(\log x + 1)}{x-1}
\end{aligned} \tag{2.24}$$

Also, note that

$$\beta_K = -\left\{\frac{r_1}{2}(\gamma + \log 4\pi) + r_2(\gamma + \log 2\pi)\right\} \leq -\frac{(\gamma + \log 2\pi)}{2}n_K < -n_K$$

The last inequality follows from $\gamma + \log 2\pi = 2.4150927\dots$ and therefore,

$$\begin{aligned}
&\frac{2\beta_K \log x}{\sqrt{x}} + \frac{n_K(\log x)(\log x + 1)}{x-1} \\
&< n_K(\log x) \left(\frac{\log x + 1}{x-1} - \frac{2}{\sqrt{x}}\right) \\
&< 0 \quad \text{for all } x \geq 3.
\end{aligned}$$

Hence we have, for all $x \geq 3$,

$$\Psi_K(x) \leq (\log x)^2 - \left(\frac{\sqrt{x}-1}{\sqrt{x}+1}\right)(\log x)(\gamma_{K,0}+1) + \frac{2\alpha_K \log x}{\sqrt{x}} \tag{2.25}$$

Estimates for $r(x)$ under GRH

Recall we wrote,

$$\begin{aligned}
r(x) &= \sum \frac{\rho(\rho-1)\log x - \rho^2 + (1-\rho)^2}{\rho^2(1-\rho)^2} \cdot \frac{x^\rho}{x-1} \\
&= \frac{\log x}{x-1} \sum \frac{x^\rho}{\rho(\rho-1)} + \frac{1}{x-1} \sum \left(\frac{x^\rho}{\rho^2} - \frac{x^\rho}{(1-\rho)^2}\right)
\end{aligned}$$

$$\begin{aligned}
&= \frac{\log x}{2(x-1)} \sum \left(\frac{x^\rho}{\rho(\rho-1)} + \frac{x^{1-\rho}}{(1-\rho)(-\rho)} \right) + \frac{1}{x-1} \sum \frac{x^{1-\rho} - x^\rho}{(1-\rho)^2} \\
&= -\frac{\log x}{2(x-1)} \sum \frac{x^\rho + x^{1-\rho}}{\rho(1-\rho)} + \frac{1}{x-1} \sum \frac{x^{1-\rho} - x^\rho}{(1-\rho)^2} \\
&= -\frac{\log x}{2(x-1)} \sum \frac{2\sqrt{x} \cos(\gamma \log x)}{\rho(1-\rho)} + \frac{1}{x-1} \sum \frac{-2i\sqrt{x} \sin(\gamma \log x)}{(1-\rho)^2}
\end{aligned}$$

where the last equality follows under GRH, from writing $\rho = \frac{1}{2} + i\gamma$ and

$$x^\rho + x^{1-\rho} = x^\rho + x^{\bar{\rho}} = \sqrt{x}e^{i\gamma \log x} + \sqrt{x}e^{-i\gamma \log x} = 2\sqrt{x} \cos(\gamma \log x)$$

etc. We also note that $|(1-\rho)^2| = \frac{1}{4} + \gamma^2 = |\rho(\rho-1)| = \rho(1-\rho)$. Thus,

$$\begin{aligned}
|r(x)| &\leq \frac{\sqrt{x} \log x}{x-1} \sum \frac{1}{|\rho(\rho-1)|} + \frac{2\sqrt{x}}{x-1} \sum \frac{1}{|(1-\rho)^2|} \\
&= \frac{\sqrt{x}(\log x + 2)}{x-1} \sum \frac{1}{\rho(1-\rho)} \\
&= \frac{2\sqrt{x}(\log x + 2)}{x-1} (\gamma_{K,0} + \alpha_K + \beta_K + 1) \quad (\text{using equation 2.4})
\end{aligned} \tag{2.26}$$

Estimates for $\ell(x)$

Recall,

$$\begin{aligned}
\ell(x) &= -\frac{r_1 + r_2}{x-1} \left[\log x + \frac{1}{2}(\log x)^2 \right] + \frac{r_1}{x-1} \left[\sum_{n=1}^{\infty} \frac{1}{(1+2n)^2} - \sum_{n=1}^{\infty} \frac{1}{4n^2} \right] - \frac{r_2}{x-1} \\
&\quad + r_1 \left[\sum_{n=1}^{\infty} \frac{2n(2n+1) \log x - 4n^2 + (1+2n)^2}{4n^2(1+2n)^2} \cdot \frac{x^{-2n}}{x-1} \right] \\
&\quad + r_2 \left[\sum_{n=1}^{\infty} \frac{n(n+1) \log x - n^2 + (1+n)^2}{n^2(1+n)^2} \cdot \frac{x^{-n}}{x-1} \right]
\end{aligned}$$

Therefore, we have, for $x \geq 3$,

$$|\ell(x)| \leq \frac{n_K(\log x)^2}{x-1} + \frac{n_K(\log x)}{x(x-1)} + \frac{2n_K}{x-1} \quad (2.27)$$

Note that, $\sum_{n=1}^{\infty} \frac{1}{n(n+1)} = 1$, whereas $\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6} \sim 1.65$.

Remark 2.4.1. Note that the above bound for $\ell(x)$ is unconditional.

We now have all the estimates to prove our theorem.

Theorem 2.4.2. Under GRH, for $|d_K| > 8$

$$|\gamma_{K,1}| \leq 2(\log \alpha_K)(2 \log \alpha_K - \gamma_{K,0}) + 18 \log \alpha_K + O\left(\frac{(\log \alpha_K)^2}{\alpha_K}\right) \quad (2.28)$$

where, as before, $\alpha_K = \log \sqrt{|d_K|}$.

Proof. Recall in (2.22) we had the formula

$$\gamma_{K,1} = \Psi_K(x) - 1 - \frac{1+x}{1-x} \log x - \frac{1}{2} \log^2 x + r(x) + \ell(x)$$

Thus,

$$|\gamma_{K,1}| \leq |\Psi_K(x)| + (\log x)^2 + |r(x)| + |\ell(x)|$$

Note that $\Psi_K(x) \geq 0$ and so $|\Psi_K(x)| = \Psi_K(x)$. Substituting the bounds obtained for $r(x)$ in (2.26), $\Psi_K(x)$ in (2.25) in the above equation we get :

$$\begin{aligned} |\gamma_{K,1}| &\leq (\log x)^2 - \left(\frac{\sqrt{x}-1}{\sqrt{x}+1}\right) (\log x)(\gamma_{K,0} + 1) + \frac{2\alpha_K \log x}{\sqrt{x}} + \\ &\quad \frac{2\sqrt{x}(\log x + 2)}{x-1} (\gamma_{K,0} + \alpha_K + \beta_K + 1) + |\ell(x)| \end{aligned}$$

Let us first focus on the coefficient α_K . We have

$$2\alpha_K \left(\frac{(\log x)}{\sqrt{x}} + \frac{\sqrt{x}(\log x + 2)}{x-1} \right) \leq \frac{10\alpha_K(\log x)}{\sqrt{x}}$$

For the coefficient of $(\gamma_{K,0} + 1)$ we have,

$$\begin{aligned}
& \frac{2\sqrt{x}(\log x + 2)}{x-1} - \left(\frac{\sqrt{x}-1}{\sqrt{x}+1} \right) (\log x) \\
&= \frac{2\sqrt{x} \log x + 4\sqrt{x} - (x+1-2\sqrt{x}) \log x}{x-1} \\
&= -\log x + \frac{4\sqrt{x} \log x + 4\sqrt{x} - 2}{x-1}
\end{aligned}$$

Now recall that $\beta_K < -n_K$, putting this together with the bounds of $\ell(x)$ from (2.27) we have

$$\begin{aligned}
& \frac{2\beta_K \sqrt{x}(\log x + 2)}{x-1} + \frac{n_K(\log x)^2}{x-1} + \frac{n_K(\log x)}{x(x-1)} + \frac{2n_K}{x-1} \\
&< \frac{n_K(\log x)}{x-1} \left[\log x + \frac{1}{x} - 2\sqrt{x} \right] + \frac{2n_K}{x-1} [1 - 2\sqrt{x}] \\
&< 0
\end{aligned}$$

Thus the cumulative contribution from terms involving β_K in $|r(x)|$ and $|\ell(x)|$ is negative. So we'll ignore these.

Therefore we have,

$$\begin{aligned}
|\gamma_{K,1}| &\leq (\log x)^2 - (\log x)(\gamma_{K,0} + 1) + \frac{10\alpha_K(\log x)}{\sqrt{x}} \\
&\quad + \left(\frac{4\sqrt{x} \log x + 4\sqrt{x} - 2}{x-1} \right) (\gamma_{K,0} + 1) \\
&\leq (\log x) \left(\log x - (\gamma_{K,0} + 1) + \frac{10\alpha_K}{\sqrt{x}} \right) \\
&\quad + \left(\frac{4\sqrt{x} \log x + 4\sqrt{x} - 2}{x-1} \right) (\gamma_{K,0} + 1)
\end{aligned}$$

Therefore, choosing $x = \alpha_K^2$ we get,

$$|\gamma_{K,1}| \leq 2(\log \alpha_K)(2 \log \alpha_K - \gamma_{K,0} + 9) + c_2 \left(\frac{(\log \alpha_K)^2}{\alpha_K} \right)$$

and hence we have our result. Note that, in the error term, the inequality follows from Ihara's upper bound on $\gamma_{K,0}$. \square

Corollary 2.4.3. When $\gamma_{K,0} \geq 0$, we get, (under GRH and for $|d_K| > 8$)

$$\gamma_{K,1} \ll \left(\log \log \sqrt{|d_K|} \right)^2$$

2.5 AN UNCONDITIONAL ARITHMETIC FORMULA FOR $\gamma_{K,1}$

Theorem 2.5.1. (Unconditionally)

$$\gamma_{K,1} = \lim_{x \rightarrow \infty} \left[\Psi_K(x) - 1 - \frac{1+x}{1-x} \log x - \frac{1}{2} (\log x)^2 \right]$$

Proof. Note that, we have the formula (as in equation 2.22)

$$\gamma_{K,1} = \Psi_K(x) - 1 - \frac{1+x}{1-x} \log x - \frac{1}{2} \log^2 x + r(x) + \ell(x)$$

Since the estimates for $\ell(x)$ in (2.27) are unconditional, we see that $\lim_{x \rightarrow \infty} \ell(x) = 0$.

To show, same is true for $r(x)$ we make use of standard zero-free region of the Dedekind zeta function. In particular, we will use the following Lemma 8.1 of [LO77] which states that :

Theorem 2.5.2. There is an absolute, effectively computable positive constant c such that $\zeta_K(s)$ has no zeros $\rho = \beta + i\gamma$ in the region :

$$|\gamma| \geq \frac{1}{1 + 4 \log d_K} , \quad \beta \geq 1 - \frac{c}{\log d_K + n_K \log(|\gamma| + 2)}$$

We have (writing $\rho = \beta + i\gamma$)

$$\begin{aligned} r(x) &= \sum \frac{\rho(\rho-1) \log x - \rho^2 + (1-\rho)^2}{\rho^2(1-\rho)^2} \cdot \frac{x^\rho}{x-1} \\ &\ll \sum \frac{(\log x) x^{\beta-1}}{\gamma^2} \end{aligned}$$

Since $\beta < 1$, we can assume the condition on β in the above theorem holds, by excluding finitely many zeros. Thus we get :

$$\begin{aligned} \sum \frac{(\log x) x^{\beta-1}}{\gamma^2} &< \sum \frac{(\log x) x^{-c(\log d_K + n_K \log(|\gamma|+2))^{-1}}}{\gamma^2} \\ &= \sum_{\log d_K + n_K \log(|\gamma|+2) < T} + \sum_{\log d_K + n_K \log(|\gamma|+2) \geq T} \end{aligned} \quad (2.29)$$

where we will choose $T = \sqrt{\log x}$. Thus for the first sum :

$$\begin{aligned} \sum_{\log d_K + n_K \log(|\gamma|+2) < T} \frac{(\log x) x^{-c(\log d_K + n_K \log(|\gamma|+2))^{-1}}}{\gamma^2} \\ < \sum \frac{(\log x) x^{-cT^{-1}}}{\gamma^2} = \left(\sum \frac{1}{\gamma^2} \right) (\log x) \exp(-c\sqrt{\log x}) \end{aligned}$$

Note that the last equality follows from :

$$\exp(-c\sqrt{\log x}) = \exp(-c \log x (\sqrt{\log x})^{-1}) = \exp(\log x^{-cT^{-1}}) = x^{-cT^{-1}}$$

Now as $x \rightarrow \infty$, clearly $\exp(-c\sqrt{\log x}) \rightarrow 0$. We also have

$$\lim_{x \rightarrow \infty} \frac{\log x}{e^{c\sqrt{\log x}}} = \lim_{y \rightarrow \infty} \frac{y^2}{e^{cy}} \rightarrow 0$$

Now let us consider the second sum in 2.29. Note that

$$\log d_K + n_K \log(|\gamma|+2) \geq \sqrt{\log x} \Rightarrow |\gamma| \geq -2 + \exp\left(\frac{\sqrt{\log x} - \log d_K}{n_K}\right)$$

We will write the expression on the right as u , i.e. $|\gamma| \geq u$. Note that as $x \rightarrow \infty$, so does u .

We will also use the following result on counting the number of zeros in a rectangle in the critical strip (it can be deduced from Jensen's theorem), see for example Theorem 5.31 of [IK21]

If $N_K(T)$ denote the number of zeros of $\zeta_K(s)$ in the region $0 \leq \Re(s) \leq 1$ and $|\Im(s)| \leq T$, then we have

$$|N_K(T+1) - N_K(T)| \ll n_K \log T + \log d_K \quad (2.30)$$

where the implied constant is absolute.

Since $x > 1$ we have

$$\begin{aligned}
& \sum_{|\gamma| \geq u} \frac{(\log x) x^{-c(\log d_K + n_K \log(|\gamma|+2))^{-1}}}{\gamma^2} \\
& < (\log x) \left(\sum_{|\gamma| \geq u} \frac{1}{\gamma^2} \right) \\
& \leq (\log x) \left(\sum_{j > u} \sum_{j < |\gamma| < j+1} \frac{1}{j^2} \right) \\
& \leq (\log x) \sum_{j > u} \frac{c_1(n_K \log j + \log d_K)}{j^2} \\
& \leq c_1 n_K (\log x) \frac{\log u + 1}{u} + c_1 (\log x) (\log d_K) \frac{1}{u} \\
& \rightarrow 0 \text{ as } x \rightarrow \infty \text{ (and therefore } u \rightarrow \infty \text{)}.
\end{aligned}$$

The above is true since,

$$\begin{aligned}
u &= -2 + \exp \left(\frac{\sqrt{\log x} - \log d_K}{n_K} \right) \\
\log(u + 2) &= \frac{\sqrt{\log x} - \log d_K}{n_K} \\
\log x &= (n_K \log(u + 2) + \log d_K)^2
\end{aligned}$$

Therefore we have, $\lim_{x \rightarrow \infty} r(x) = 0$ and this completes the proof. \square

In the next section we will show that our computations and techniques generalize, and prove similar results for the m -th Euler-Kronecker constants $\gamma_{K,m}$.

2.6 GENERALIZATIONS TO HIGHER CONSTANTS $\gamma_{K,m}$

Recall, that we wrote the Laurent series of $\frac{\zeta'_K(s)}{\zeta_K(s)}$ about $s = 1$, to be :

$$\begin{aligned} \frac{\zeta'_K(s)}{\zeta_K(s)} &= \frac{-1}{s-1} + \sum_{m=0}^{\infty} \gamma_{K,m} (s-1)^m \\ \Rightarrow \frac{\zeta'_K(s)}{\zeta_K(s)} + \frac{1}{s-1} &= \sum_{m=0}^{\infty} \gamma_{K,m} (s-1)^m \end{aligned}$$

Differentiating both sides m times and letting $s \rightarrow 1$ we get,

$$\lim_{s \rightarrow 1} \frac{d^m}{ds^m} \left[\frac{\zeta'_K(s)}{\zeta_K(s)} + \frac{1}{s-1} \right] = m! \cdot \gamma_{K,m} \quad (2.31)$$

On the other hand, from the Hadamard factorization, we had

$$\begin{aligned} -\frac{\zeta'_K(s)}{\zeta_K(s)} &= \frac{1}{s} + \frac{1}{s-1} - \sum \frac{1}{s-\rho} + \alpha_K + \beta_K + \tilde{\Gamma}_K(s) \\ \Rightarrow \frac{\zeta'_K(s)}{\zeta_K(s)} + \frac{1}{s-1} &= -\frac{1}{s} + \sum \frac{1}{s-\rho} - \alpha_K - \beta_K - \tilde{\Gamma}_K(s) \end{aligned}$$

So, differentiating both sides of the above m times we have,

$$\frac{d^m}{ds^m} \left[\frac{\zeta'_K(s)}{\zeta_K(s)} + \frac{1}{s-1} \right] = -\frac{(-1)^m m!}{s^{m+1}} + \sum \frac{(-1)^m m!}{(s-\rho)^{m+1}} - \tilde{\Gamma}_K^{(m)}(s) \quad (2.32)$$

Letting $s \rightarrow 1$ in the above and using equation (2.31), we derive the formula

$$\begin{aligned} m! \cdot \gamma_{K,m} &= -(-1)^m m! + \sum \frac{(-1)^m m!}{(1-\rho)^{m+1}} - \tilde{\Gamma}_K^{(m)}(1) \\ \Rightarrow \boxed{\gamma_{K,m} + (-1)^m} &= \sum \frac{(-1)^m}{(1-\rho)^{m+1}} - \frac{1}{m!} \tilde{\Gamma}_K^{(m)}(1) \end{aligned} \quad (2.33)$$

Rewriting equation (2.32) to match our $m = 1$ setting :

$$\frac{d^m}{ds^m} \frac{\zeta'_K(s)}{\zeta_K(s)} = -\frac{(-1)^m m!}{(s-1)^{m+1}} - \frac{(-1)^m m!}{s^{m+1}} + \sum \frac{(-1)^m m!}{(s-\rho)^{m+1}} - \tilde{\Gamma}_K^{(m)}(s) \quad (2.34)$$

Recall, we were writing,

$$Z_K(s) = -\frac{\zeta'_K(s)}{\zeta_K(s)}$$

So that, equation (2.34) above, takes the shape : (note the change of signs)

$$Z_K^{(m)}(s) = \frac{(-1)^m m!}{(s-1)^{m+1}} + \frac{(-1)^m m!}{s^{m+1}} + \sum \frac{(-1)^{m+1} m!}{(s-\rho)^{m+1}} + \tilde{\Gamma}_K^{(m)}(s) \quad (2.35)$$

On the other hand, from the Euler product, we had :

$$Z_K(s) = -\frac{\zeta'_K(s)}{\zeta_K(s)} = \sum_{P, k \geq 1} \frac{\log N(P)}{N(P)^{ks}}$$

Differentiating m times yields

$$Z_K^{(m)}(s) = \sum_{P, k \geq 1} \frac{(-1)^m k^m (\log N(P))^{m+1}}{N(P)^{ks}} \quad (2.36)$$

To deduce bounds and an exact formula, we follow the same steps as the $m = 1$ case, and evaluate the integral

$$\Psi_K^{(\mu)}(m, x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^{s-\mu}}{s-\mu} Z_K^{(m)}(s) ds \quad \text{for } c \gg 1$$

We will compute the expression $x\Psi_K^{(1)}(m, x) - \Psi_K^{(0)}(m, x)$, i.e. for $\mu = 0$ and 1, in two different ways using equation (2.35) and (2.36). As a result, on one side, the constituents of $\gamma_{K,m}$ as in equation (2.33), will occur. Whereas, on the other side we will have an expression of x and choosing appropriate x will lead us to bounds on these higher coefficients.

The following classical formula will come in handy.

$$\frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{y^s}{s} ds = \begin{cases} 0 & 0 < y < 1 \\ \frac{1}{2} & y = 1 \\ 1 & y > 1 \end{cases} \quad (2.37)$$

First using equation (2.36), we get

$$\begin{aligned}
 & x\Psi_K^{(1)}(m, x) - \Psi_K^{(0)}(m, x) \\
 &= x \cdot \sum_{P, k \geq 1} \frac{(-1)^m k^m (\log N(P))^{m+1}}{N(P)^k} \left[\frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{1}{s-1} \left(\frac{x}{N(P)^k} \right)^{s-1} ds \right] \\
 &\quad - \sum_{P, k \geq 1} (-1)^m k^m (\log N(P))^{m+1} \left[\frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{1}{s} \left(\frac{x}{N(P)^k} \right)^s ds \right] \\
 &= x \left[\sum_{k, N(P)^k < x} \frac{(-1)^m k^m (\log N(P))^{m+1}}{N(P)^k} + \sum_{k, N(P)^k = x} \frac{(-1)^m k^m (\log N(P))^{m+1}}{N(P)^k} \cdot \frac{1}{2} \right] \\
 &\quad - \left[\sum_{k, N(P)^k < x} (-1)^m k^m (\log N(P))^{m+1} + \sum_{k, N(P)^k = x} (-1)^m k^m (\log N(P))^{m+1} \cdot \frac{1}{2} \right] \\
 &= \sum_{k, N(P)^k < x} \left(\frac{x}{N(P)^k} - 1 \right) (-1)^m k^m (\log N(P))^{m+1}
 \end{aligned}$$

Note that the second sums in the square brackets, cancel each other as $N(P)^k = x$ and there is an x at the very beginning of that line. We define :

$$\boxed{\Psi_K(m, x) = \frac{1}{x-1} \sum_{k, N(P)^k \leq x} \left(\frac{x}{N(P)^k} - 1 \right) k^m (\log N(P))^{m+1} \quad \text{for } x > 1} \tag{2.38}$$

Therefore, on one side we have,

$$x\Psi_K^{(1)}(m, x) - \Psi_K^{(0)}(m, x) = (-1)^m (x-1) \Psi_K(m, x) \tag{2.39}$$

We now focus on equation (2.35) and first look at the contribution from the term : $(-1)^m m! \left[\frac{1}{s^{m+1}} + \frac{1}{(s-1)^{m+1}} \right]$. To evaluate this, we will use a generalization of the classical formula stated in equation (2.37).

For $n \geq 1$

$$\frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{y^s}{s^{n+1}} ds = \begin{cases} 0 & 0 < y \leq 1 \\ \frac{1}{n!} (\log y)^n & y > 1 \end{cases} \tag{2.40}$$

Let us write,

$$f(m, x) = \frac{(-1)^m m!}{2\pi i} \int_{(c)} x^s \left[\frac{1}{s-1} \left(\frac{1}{s^{m+1}} + \frac{1}{(s-1)^{m+1}} \right) - \frac{1}{s} \left(\frac{1}{s^{m+1}} + \frac{1}{(s-1)^{m+1}} \right) \right] ds$$

This will be the net contribution coming from the first two terms of (2.35). We will simplify this a little bit and write a recursive relation satisfied by $f(m, x)$, which will in turn help us to estimate it.

$$\begin{aligned} f(m, x) &= \frac{(-1)^m m!}{2\pi i} \int_{(c)} x^s \left[\frac{1}{s^{m+1}(s-1)} + \frac{1}{(s-1)^{m+2}} - \frac{1}{s^{m+2}} - \frac{1}{s(s-1)^{m+1}} \right] ds \\ &= (-1)^m m! \left[x \frac{(\log x)^{m+1}}{(m+1)!} - \frac{(\log x)^{m+1}}{(m+1)!} \right] + \\ &\quad \frac{(-1)^m m!}{2\pi i} \int_{(c)} x^s \left[\frac{1}{s^{m+1}(s-1)} - \frac{1}{s(s-1)^{m+1}} \right] ds \\ &= \frac{(-1)^m}{m+1} (x-1)(\log x)^{m+1} + \\ &\quad \frac{(-1)^m m!}{2\pi i} \int_{(c)} x^s \left[\frac{1}{s^m(s-1)} - \frac{1}{s^{m+1}} + \frac{1}{s(s-1)^m} - \frac{1}{(s-1)^{m+1}} \right] ds \\ &= \frac{(-1)^m}{m+1} (x-1)(\log x)^{m+1} + (-1)^{m+1} (x+1)(\log x)^m + \\ &\quad \frac{(-1)^m m!}{2\pi i} \int_{(c)} x^s \left[\frac{1}{s^m(s-1)} + \frac{1}{s(s-1)^m} \right] ds \\ &= \frac{(-1)^m}{m+1} (x-1)(\log x)^{m+1} + (-1)^{m+1} (x+1)(\log x)^m + \\ &\quad \frac{(-1)^m (m)!}{2\pi i} \int_{(c)} x^s \left[\frac{1}{s^{m-1}(s-1)} - \frac{1}{s^m} - \frac{1}{s(s-1)^{m-1}} + \frac{1}{(s-1)^m} \right] ds \\ &= \frac{(-1)^m}{m+1} (x-1)(\log x)^{m+1} + (-1)^{m+1} (x+1)(\log x)^m + m(m-1)f(m-2, x) \end{aligned} \tag{2.41}$$

Note that from our previous computations, e.g. see equation (2.16) we get the $m = 1$ case and from [Ihao6], 1.3.15 for the $m = 0$ case :

$$\begin{aligned} f(1, x) &= (1-x) \left[2 + \frac{1+x}{1-x} \log x + \frac{1}{2} (\log x)^2 \right] \\ f(0, x) &= (x-1) \log(x) \end{aligned}$$

Now let us compute the contribution from the sum of non-trivial zeros. We are trying to evaluate :

$$\frac{(-1)^{m+1} m!}{2\pi i} \int_{c-i\infty}^{c+i\infty} \sum x^s \left[\frac{1}{(s-1)(s-\rho)^{m+1}} - \frac{1}{s(s-\rho)^{m+1}} \right] ds \quad (2.42)$$

Similar to the $m = 1$ case we do this by contour manipulation. We can take a rectangular contour as in Figure 2.1, and show that as $T \rightarrow \infty$ the value of the integral goes to zero on each side of the rectangle, except $c - iT \rightarrow c + iT$. Thus, by residue theorem, the line integral on (c) will be equal to the sum of residues.

For the second term in the square brackets above, in (2.42) :

the pole at $s = 0$ has residue : $(-1)^{m+2} m! \cdot \frac{1}{(-\rho)^{m+1}} = -\frac{m!}{\rho^{m+1}}$.

The pole at ρ (order = $m + 1$) has residue :

$$\begin{aligned} &(-1)^{m+2} m! \lim_{s \rightarrow \rho} \frac{d^m}{ds^m} \left(\frac{x^s}{s} \right) \\ &= (-1)^{m+2} m! \lim_{s \rightarrow \rho} \sum_{k=0}^m \binom{m}{k} \frac{d^{(m-k)}}{ds^{(m-k)}} x^s \cdot \frac{d^k}{ds^k} \frac{1}{s} \\ &= (-1)^{m+2} m! \lim_{s \rightarrow \rho} \sum_{k=0}^m \binom{m}{k} x^s (\log x)^{m-k} \frac{(-1)^k k!}{s^{k+1}} \\ &= (-1)^{m+2} m! \sum_{k=0}^m \binom{m}{k} x^\rho (\log x)^{m-k} \frac{(-1)^k k!}{\rho^{k+1}} \end{aligned}$$

Computations for the first term are very similar.

Residue at $s = 1$ will be $(-1)^{m+1}m!x \sum \frac{1}{(1-\rho)^{m+1}}$.

The pole at ρ (order = $m + 1$) has residue :

$$\begin{aligned} & (-1)^{m+1}m! \lim_{s \rightarrow \rho} \frac{d^m}{ds^m} \left(\frac{x^s}{s-1} \right) \\ &= (-1)^{m+1}m! \lim_{s \rightarrow \rho} \sum_{k=0}^m \binom{m}{k} \frac{d^{(m-k)}}{ds^{(m-k)}} x^s \cdot \frac{d^k}{ds^k} \frac{1}{s-1} \\ &= (-1)^{m+1}m! \sum_{k=0}^m \binom{m}{k} x^\rho (\log x)^{m-k} \frac{(-1)^k k!}{(\rho-1)^{k+1}} \end{aligned}$$

Note that for $m = 1$ case, the residues from $s = 0$ and $s = 1$ came together to form a constituent of $\gamma_{K,1}$, whereas the rest of the terms involving x , we wrote it as $r(x)$ and estimated it. We will do the same here, but we need to be a bit more careful. Since the power of $(-1)^m$ gets canceled in computing the pole at 0. We write the net contribution as follows :

$$m!(-1)^{m+1}(x-1) \left[\sum \frac{1}{(1-\rho)^{m+1}} + r(m,x) \right] \quad (2.43)$$

where

$$\begin{aligned} r(m,x) &= \frac{1}{x-1} \left[((-1)^{m+1} - 1) \sum \frac{1}{\rho^{m+1}} \right] + \\ & \sum_{\rho} \left(\sum_{k=0}^m (-1)^k k! \binom{m}{k} \left[\frac{1}{(\rho-1)^{k+1}} - \frac{1}{\rho^{k+1}} \right] \frac{x^\rho (\log x)^{m-k}}{x-1} \right) \end{aligned}$$

Now let us compute the contribution from the Gamma factors. Recall,

$$\tilde{\Gamma}_K(s) = \frac{r_1}{2} \left[\psi \left(\frac{s}{2} \right) - \psi \left(\frac{1}{2} \right) \right] + r_2 [\psi(s) - \psi(1)]$$

where $\psi(s) = \frac{\Gamma'}{\Gamma}(s)$ is the *digamma* function. Thus, taking the m -th derivative we get,

$$\tilde{\Gamma}_K^{(m)}(s) = \frac{r_1}{2^{m+1}} \psi^{(m)} \left(\frac{s}{2} \right) + r_2 \psi^{(m)}(s)$$

We look at the series expansion of $\frac{\Gamma'}{\Gamma}(s)$ and its m -th derivative.

$$\begin{aligned}\frac{\Gamma'}{\Gamma}(s) &= \psi(s) = -\gamma - \sum_{k=0}^{\infty} \left(\frac{1}{s+k} - \frac{1}{1+k} \right) \\ \Rightarrow \psi^{(m)}(s) &= - \sum_{k=0}^{\infty} \frac{(-1)^m m!}{(s+k)^{m+1}}\end{aligned}$$

Note that the above series expansion of digamma is valid in the entire complex plane except for non-positive integers. i.e. in $\mathbb{C} \setminus \{0, -1, -2, -3, \dots\}$.

Therefore we have the following expression for $\tilde{\Gamma}_K^{(m)}(s)$:

$$\begin{aligned}\tilde{\Gamma}_K^{(m)}(s) &= -\frac{r_1}{2^{m+1}} \sum_{k=0}^{\infty} \frac{(-1)^m m! 2^{m+1}}{(s+2k)^{m+1}} - r_2 \sum_{k=0}^{\infty} \frac{(-1)^m m!}{(s+k)^{m+1}} \\ &= (-1)^{m+1} m! \left[\frac{(r_1+r_2)}{s^{m+1}} + r_1 \sum_{k=1}^{\infty} \frac{1}{(s+2k)^{m+1}} + r_2 \sum_{k=1}^{\infty} \frac{1}{(s+k)^{m+1}} \right]\end{aligned}\tag{2.44}$$

In the last line, we have separated the $k = 0$ case, and the sums now start from $k = 1$. The summand $k = 0$ will give us the main term in the contribution from these Gamma factors.

Remember, we are trying to evaluate the integral

$$\frac{x}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^{s-1}}{s-1} \tilde{\Gamma}_K^{(m)}(s) ds - \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^s}{s} \tilde{\Gamma}_K^{(m)}(s) ds$$

We first compute the contribution from $\frac{(r_1+r_2)}{s^{m+1}}$ term. We have the partial fraction decomposition :

$$\frac{1}{s^{m+1}(s-1)} = \frac{1}{s-1} - \frac{1}{s} - \frac{1}{s^2} - \dots - \frac{1}{s^{m+1}}$$

So contribution from the first part looks like :

$$(-1)^{m+1} m! (r_1+r_2) \left[x-1 - \sum_{k=1}^{m+1} \frac{1}{(k)!} (\log x)^k \right]\tag{2.45}$$

We have used the classical formula (2.40) multiple times to get the above. Now to compute the contribution from the series terms in (2.44), we first notice the similarity of it to the sum on non-trivial zeros $\sum \frac{(-1)^m m!}{(s-\rho)^{m+1}}$. So, the residue computations will be very similar, only ρ replaced by $-2k$ or $-k$. For the first series, we are looking at the integral :

$$(-1)^{m+1} m! r_1 \left[\frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} x^s \left(\sum_{k=1}^{\infty} \frac{1}{(s-1)(s+2k)^{m+1}} - \frac{1}{s(s+2k)^{m+1}} \right) ds \right]$$

In the second term,

the pole at $s = 0$ has residue : $\frac{(-1)^{m+2} m! r_1}{(2k)^{m+1}}$.

The pole at $s = -2k$ (order = $m + 1$) has residue :

$$\begin{aligned} & (-1)^{m+2} m! r_1 \lim_{s \rightarrow (-2k)} \frac{d^m}{ds^m} \left(\frac{x^s}{s} \right) \\ &= (-1)^{m+2} m! r_1 \lim_{s \rightarrow (-2k)} \sum_{j=0}^m \binom{m}{j} \frac{d^{(m-j)}}{ds^{(m-j)}} x^s \cdot \frac{d^j}{ds^j} \frac{1}{s} \\ &= (-1)^{m+2} m! r_1 \sum_{j=0}^m \binom{m}{j} x^{-2k} (\log x)^{m-j} \frac{(-1)^j j!}{(-2k)^{j+1}} \end{aligned}$$

Computations for the first term are very similar.

Residue at $s = 1$ will be $\frac{(-1)^{m+1} m! r_1}{(1+2k)^{m+1}} x$.

The pole at $s = -2k$ (order = $m + 1$) has residue :

$$\begin{aligned} & (-1)^{m+1} m! r_1 \lim_{s \rightarrow (-2k)} \frac{d^m}{ds^m} \left(\frac{x^s}{s-1} \right) \\ &= (-1)^{m+1} m! r_1 \lim_{s \rightarrow (-2k)} \sum_{j=0}^m \binom{m}{j} \frac{d^{(m-j)}}{ds^{(m-j)}} x^s \cdot \frac{d^j}{ds^j} \frac{1}{s-1} \\ &= (-1)^{m+1} m! r_1 \sum_{j=0}^m \binom{m}{j} x^{-2k} (\log x)^{m-j} \frac{(-1)^j j!}{(-2k-1)^{j+1}} \end{aligned}$$

Thus the net contribution from the first series of equation (2.44)

$$\begin{aligned}
 & (-1)^{m+1} m! r_1 \left[x \sum_{k=1}^{\infty} \frac{1}{(1+2k)^{m+1}} - \sum_{k=1}^{\infty} \frac{1}{(2k)^{m+1}} \right] + \\
 & (-1)^{m+1} m! r_1 \sum_{k=1}^{\infty} \left[\sum_{j=0}^m \binom{m}{j} x^{-2k} (\log x)^{m-j} j! \left(\frac{1}{(2k)^{j+1}} - \frac{1}{(2k+1)^{j+1}} \right) \right]
 \end{aligned} \tag{2.46}$$

Looking at this, we can precisely write down the contribution from the second series of (2.44), by just replacing $2k$ by k and r_1 by r_2 . We get,

$$\begin{aligned}
 & (-1)^{m+1} m! r_2 \left[x \sum_{k=1}^{\infty} \frac{1}{(1+k)^{m+1}} - \sum_{k=1}^{\infty} \frac{1}{(k)^{m+1}} \right] + \\
 & (-1)^{m+1} m! r_2 \sum_{k=1}^{\infty} \left[\sum_{j=0}^m \binom{m}{j} x^{-k} (\log x)^{m-j} j! \left(\frac{1}{(k)^{j+1}} - \frac{1}{(k+1)^{j+1}} \right) \right] \\
 & = (-1)^{m+1} m! r_2 (x-1) \sum_{k=1}^{\infty} \frac{1}{(1+k)^{m+1}} + (-1)^m m! r_2 + \\
 & (-1)^{m+1} m! r_2 \sum_{k=1}^{\infty} \left[\sum_{j=0}^m \binom{m}{j} x^{-k} (\log x)^{m-j} j! \left(\frac{1}{(k)^{j+1}} - \frac{1}{(k+1)^{j+1}} \right) \right]
 \end{aligned} \tag{2.47}$$

We will summarize these computations in equations (2.45), (2.46) and (2.47) to write the net contribution in the following form :

$$(x-1) \left[\tilde{\Gamma}_K^{(m)}(1) + (-1)^{m+1} m! \ell(m, x) \right] \tag{2.48}$$

where

$$\begin{aligned} \ell(m, x) &= \frac{r_1 + r_2}{(x-1)} \sum_{t=1}^{m+1} \frac{(\log x)^t}{t!} + \frac{r_1}{(x-1)} \sum_{k=1}^{\infty} \left[\frac{1}{(1+2k)^{m+1}} - \frac{1}{(2k)^{m+1}} \right] \\ &\quad - \frac{r_2}{(x-1)} + \frac{r_1}{(x-1)} \sum_{k=1}^{\infty} \left[\sum_{j=0}^m \binom{m}{j} x^{-2k} (\log x)^{m-j} j! \left(\frac{1}{(2k)^{j+1}} - \frac{1}{(2k+1)^{j+1}} \right) \right] \\ &\quad + \frac{r_2}{(x-1)} \sum_{k=1}^{\infty} \left[\sum_{j=0}^m \binom{m}{j} x^{-k} (\log x)^{m-j} j! \left(\frac{1}{(k)^{j+1}} - \frac{1}{(k+1)^{j+1}} \right) \right] \end{aligned} \tag{2.49}$$

2.7 BOUNDS FOR $\gamma_{K,m}$ UNDER GRH

Let us first estimate $\ell(m, x)$ as the expression is right above.

Lemma 2.7.1. For $m \geq 1$, we have

$$|\ell(m, x)| \leq \frac{n_K}{x-1} \left(e (\log x)^{m+1} + 1 + \frac{(m+1)! (\log x)^m}{x} \right)$$

where n_K denotes the degree $[K : \mathbb{Q}]$. Also, $n_K = r_1 + 2r_2$.

Proof. Note that after taking absolute value and applying some triangle inequalities in (2.49), the first sum is $\leq \frac{en_K (\log x)^{m+1}}{x-1}$. The next two terms combined is $\leq \frac{n_K}{x-1}$.

Now let us have a closer look at the series terms.

As both $x, k > 1$, we have $x^{-k} \leq x^{-1}$. We'll also write, $\binom{m}{j} j! \leq m!$.

Thus we have,

$$\begin{aligned} &\sum_{j=0}^m \binom{m}{j} x^{-k} (\log x)^{m-j} j! \left(\frac{1}{(k)^{j+1}} - \frac{1}{(k+1)^{j+1}} \right) \\ &\leq \frac{m! (\log x)^m}{x} \sum_{j=0}^m \left(\frac{1}{(k)^{j+1}} - \frac{1}{(k+1)^{j+1}} \right) \\ &\leq \frac{(m+1)! (\log x)^m}{x} \frac{1}{k(k+1)} \end{aligned}$$

Therefore we have the result! □

Remark 2.7.2. Note that we didn't need GRH for the above estimate. Lemma 2.7.1 is unconditional.

Let us now estimate, $\Psi_K(m, x)$. First note that it is always non-negative.

Lemma 2.7.3. For all $x \geq 3$, we have

$$\Psi_K(m, x) \leq (\log x)^m \left(\log x - \frac{\sqrt{x}-1}{\sqrt{x}+1}(\gamma_{K,0} + 1) + \frac{2\alpha_K}{\sqrt{x}} \right) \quad (2.50)$$

Proof.

$$\begin{aligned} \Psi_K(m, x) &= \frac{1}{x-1} \sum_{k, N(P)^k \leq x} \left(\frac{x}{N(P)^k} - 1 \right) k^m (\log N(P))^{m+1} \\ &= \frac{1}{x-1} \sum_{k, N(P)^k \leq x} \left(\frac{x}{N(P)^k} - 1 \right) k (\log N(P))^2 (\log N(P)^k)^{m-1} \\ &\leq (\log x)^{m-1} \Psi_K(x) \quad [\text{where } \Psi_K(x) = \Psi_K(1, x) \text{ as in (2.14)}] \\ &\leq (\log x)^m \left(\log x - \frac{\sqrt{x}-1}{\sqrt{x}+1}(\gamma_{K,0} + 1) + \frac{2\alpha_K}{\sqrt{x}} \right) \end{aligned}$$

where the last inequality is for all $x \geq 3$ and follows from (2.25). \square

Lemma 2.7.4. For $m \geq 1$, under GRH

$$|r(m, x)| \leq \frac{(m!2^{m+1}\sqrt{x}(\log x)^m + 4)}{x-1} (\gamma_{K,0} + \alpha_K + \beta_K + 1)$$

Proof. Recall,

$$\begin{aligned} r(m, x) &= \frac{1}{x-1} \left[((-1)^{m+1} - 1) \sum \frac{1}{\rho^{m+1}} \right] + \\ &\quad \sum_{\rho} \left(\sum_{k=0}^m (-1)^k k! \binom{m}{k} \left[\frac{1}{(\rho-1)^{k+1}} - \frac{1}{\rho^{k+1}} \right] \frac{x^\rho (\log x)^{m-k}}{x-1} \right) \end{aligned} \quad (2.51)$$

Note that, for m odd, the first sum of $r(m, x)$ above, is zero. This is why, such a term didn't not appear in our computations for $\gamma_{K,1}$.

For $m \geq 2$, m even,

$$\left| \frac{2}{x-1} \sum \frac{1}{\rho^{m+1}} \right| \leq \frac{2}{x-1} \sum \frac{1}{|\rho|^{m+1}} \leq \frac{2}{x-1} \sum \frac{1}{|\rho|^2}$$

As in the $m = 1$ case, under GRH we can write $\rho = \frac{1}{2} + i\gamma$ and therefore, $|\rho|^2 = \frac{1}{4} + \gamma^2 = \rho(1-\rho)$ and so,

$$\sum \frac{1}{|\rho|^2} = \sum \frac{1}{\rho(1-\rho)} = 2(\gamma_{K,0} + \alpha_K + \beta_K + 1)$$

e.g. see equation (2.4).

Now for the second sum,

$$\begin{aligned} & \sum_{\rho} \left| \sum_{k=0}^m (-1)^k k! \binom{m}{k} \left[\frac{1}{(\rho-1)^{k+1}} - \frac{1}{\rho^{k+1}} \right] \frac{x^{\rho} (\log x)^{m-k}}{x-1} \right| \\ & \leq m! \frac{\sqrt{x} (\log x)^m 2^m}{x-1} \sum_{\rho} \frac{1}{|\rho(1-\rho)|} \\ & \leq (m! 2^{m+1}) \frac{\sqrt{x} (\log x)^m}{x-1} (\gamma_{K,0} + \alpha_K + \beta_K + 1) \end{aligned}$$

The last two inequalities are under GRH and follows from the same reasoning given right above. \square

Note that, we have kept the β_K here on purpose. It is negative (in fact, as we showed before, $\beta_K < -n_K$) and it will help us ignore most, if not all, of the contribution coming from the gamma factors, as we shall see shortly.

We are now ready to state and prove our results on bounds for the general n -th Euler-Kronecker constants :

Theorem 2.7.5. Under GRH, for $d_K \geq 8$, and $m \geq 1$

$$\gamma_{K,m} \ll \frac{2^m}{m!} (\log(2^m (m!)^2) + \log \alpha_K)^m (2 \log(2^m (m!)^2) + 2 \log \alpha_K - \gamma_{K,0} + 1)$$

Proof. Putting our above computations together, the left hand side, from equation (2.39) is : $(-1)^m(x-1)\Psi_K(m, x)$. Whereas, the right hand side is obtained from combining equations (2.41), (2.43) and (2.48). We have,

$$\begin{aligned} (-1)^m(x-1)\Psi_K(m, x) &= f(m, x) + m!(-1)^{m+1}(x-1) \left[\sum \frac{1}{(1-\rho)^{m+1}} + r(m, x) \right] \\ &\quad + (x-1) \left[\tilde{\Gamma}_K^{(m)}(1) + (-1)^{m+1}m! \ell(m, x) \right] \\ \Rightarrow (-1)^m\Psi_K(m, x) &= \frac{f(m, x)}{(x-1)} - \left[\sum \frac{(-1)^m m!}{(1-\rho)^{m+1}} - \tilde{\Gamma}_K^{(m)}(1) \right] + \\ &\quad m!(-1)^{m+1} [r(m, x) + \ell(m, x)] \end{aligned}$$

Now from equation (2.33) we get :

$$\begin{aligned} \Rightarrow (-1)^m\Psi_K(m, x) &= \frac{f(m, x)}{(x-1)} - m! [\gamma_{K,m} + (-1)^m] + \\ &\quad m!(-1)^{m+1} [r(m, x) + \ell(m, x)] \end{aligned}$$

$$m! [\gamma_{K,m} + (-1)^m] = (-1)^{m+1}\Psi_K(m, x) + \frac{f(m, x)}{(x-1)} + (-1)^{m+1}m! [r(m, x) + \ell(m, x)] \quad (2.52)$$

Note that, as in the $m = 1$ case, it is easily checked that the cumulative contribution from the β_K term in Lemma 2.7.4 and $|\ell(m, x)|$ is negative. And by Ihara's bound, $\gamma_{K,0}$ is dominated by α_K . Therefore, we can write

$$r(m, x) + \ell(m, x) \ll \frac{m!2^{m+1}(\log x)^m \alpha_K}{\sqrt{x}}$$

Note that the term involving α_K in $\Psi_K(m, x)$ as in Lemma 2.7.3 gets absorbed in this as well. Also we see from (2.41) that $|\frac{f(m, x)}{x-1}| \ll (\log x)^{m+1}$

Putting this together with Lemma 2.7.3 we get

$$m![\gamma_{K,m} + (-1)^m] \ll (\log x)^m \left(\log x - (\gamma_{K,0} + 1) + \frac{(m!)^2 2^{m+1} \alpha_K}{\sqrt{x}} \right)$$

Choosing $x = ((m!)^2 2^m \alpha_K)^2$ minimizes the second term in brackets and gives us the bound :

$$\gamma_{K,m} \ll \frac{2^m}{m!} (\log(2^m (m!)^2) + \log \alpha_K)^m (2 \log(2^m (m!)^2) + 2 \log \alpha_K - \gamma_{K,0} + 1)$$

□

Corollary 2.7.6. If $\gamma_{K,0} \geq 0$ together with the conditions of the theorem, we get

$$\gamma_{K,m} \ll \frac{2^m}{m!} (\log(2^m (m!)^2) + \log \alpha_K)^{m+1}$$

In particular, for $m \ll \frac{\log d_K}{\log \log d_K}$, we have

$$\gamma_{K,m} \ll \frac{2^{m+1}}{m!} (\log \log d_K)^{m+1}$$

2.8 AN UNCONDITIONAL ARITHMETIC FORMULA FOR $\gamma_{K,m}$

We can also prove the following general arithmetic formula like that in Theorem 2.5.1, unconditionally.

Theorem 2.8.1.

$$\gamma_{K,m} + (-1)^m = \lim_{x \rightarrow \infty} \frac{1}{m!} \left[(-1)^{m+1} \Psi_K(m, x) + \frac{f(m, x)}{(x-1)} \right] \quad (2.53)$$

where as before $f(m, x)$ is recursively defined as :

$$f(m, x) = \frac{(-1)^m}{m+1} (x-1) (\log x)^{m+1} + (-1)^{m+1} (x+1) (\log x)^m + m(m-1) f(m-2, x)$$

$$f(1, x) = (1-x) \left[2 + \frac{1+x}{1-x} \log x + \frac{1}{2} (\log x)^2 \right]$$

$$f(0, x) = (x-1) \log(x)$$

and

$$\Psi_K(m, x) = \frac{1}{x-1} \sum_{k, N(P)^k \leq x} \left(\frac{x}{N(P)^k} - 1 \right) k^m (\log N(P))^{m+1} \quad \text{for } x > 1$$

Proof. As we saw in proof of Theorem 2.7.5, equation (2.52) :

$$m! [\gamma_{K,m} + (-1)^m] = (-1)^{m+1} \Psi_K(m, x) + \frac{f(m, x)}{(x-1)} + (-1)^{m+1} m! [r(m, x) + \ell(m, x)]$$

From Lemma (2.7.1) we get $\lim_{x \rightarrow \infty} \ell(m, x) = 0$. We will show the same for $r(m, x)$ as well, which will give us the result.

Recall, it was the $r(m, x)$ term which we estimated using GRH. To prove an unconditional result, we will reduce it to the case of the proof of Theorem 2.5.1. Recall, we wrote

$$r(m, x) = \frac{1}{x-1} \left[((-1)^{m+1} - 1) \sum \frac{1}{\rho^{m+1}} \right] + \sum_{\rho} \left(\sum_{k=0}^m (-1)^k k! \binom{m}{k} \left[\frac{1}{(\rho-1)^{k+1}} - \frac{1}{\rho^{k+1}} \right] \frac{x^{\rho} (\log x)^{m-k}}{x-1} \right)$$

For fixed m , note that the first term goes to 0 as $x \rightarrow \infty$. Writing $\rho = \beta + i\gamma$, we get

$$r(m, x) \ll 2^m \cdot m! (\log x)^m \sum \frac{x^{\beta-1}}{\gamma^2}$$

We note that in the computations for the proof of Theorem 2.5.1, if we replace $\log x$ by $(\log x)^m$, it still works, giving us $\lim_{x \rightarrow \infty} r(m, x) = 0$. Therefore, we have

$$\gamma_{K,m} + (-1)^m = \lim_{x \rightarrow \infty} \frac{1}{m!} \left[(-1)^{m+1} \Psi_K(m, x) + \frac{f(m, x)}{(x-1)} \right]$$

□

BOUNDS FOR THE FUNCTION FIELD CASE

3.1 PRELIMINARIES

In this chapter we deduce similar bounds as in Chapter 2 for the function field case. Let q be a power of a prime and \mathbb{F}_q be the finite field with q elements. Let K be the function field of a curve X over \mathbb{F}_q of genus g . A good reference for the basic facts about the zeta function $\zeta_K(s)$ is [Ros00].

We set $u = q^{-s}$, then the $\zeta_K(s)$ is a rational function of u of the form

$$\zeta_K(s) = \frac{\prod_{i=1}^g (1 - \pi_i u)(1 - \bar{\pi}_i u)}{(1 - u)(1 - qu)} \quad \text{with } \pi_i \bar{\pi}_i = q \quad \text{for all } 1 \leq i \leq g \quad (3.1)$$

Note that each zero $\frac{1}{\pi_i}$ or $\frac{1}{\bar{\pi}_i}$ of $\zeta_K(s)$ in u corresponds to infinitely many zeros in s and all of them are translations of a zero by $\frac{2\pi i n}{\log q}$, $n \in \mathbb{Z}$. Similarly, poles are translations of 0 and 1 by $\frac{2\pi i n}{\log q}$, $n \in \mathbb{Z}$.

Also, $\zeta_K(s)$ has a simple pole at $s = 1$, and thus like the number field case we can write the Laurent series of it's logarithmic derivative as

$$\frac{\zeta'_K(s)}{\zeta_K(s)} = \frac{-1}{s-1} + \gamma_{K,0} + \gamma_{K,1}(s-1) + \dots \quad (3.2)$$

and define $\gamma_{K,m}$ as the general m -th *Euler-Kronecker* constants.

Ihara in [Iha06], 1.3.10 derives the following Stark like lemma.

Lemma 3.1.1.

$$-\frac{\zeta'_K(s)}{\zeta_K(s)} = \frac{1}{s} + \frac{1}{s-1} - \sum_{\rho} \frac{1}{s-\rho} + (g-1) \log q + \sum_{\theta \neq 0,1} \frac{1}{s-\theta} \quad (3.3)$$

where ρ runs over the non-trivial zeros of and θ runs over all poles $\neq 0, 1$ of $\zeta_K(s)$.

Proof. Write a simpler rational form $\zeta_K(s) = \prod_{\alpha \in A} (1 - \alpha q^{-s})^{\lambda_\alpha}$, where $\lambda_\alpha = \pm 1$ and A is a finite subset of \mathbb{C}^\times . Taking the logarithmic derivative we get,

$$\frac{\zeta'_K(s)}{\zeta_K(s)} = \sum_{\alpha \in A} \lambda_\alpha \frac{-\alpha q^{-s} \log q}{1 - \alpha q^{-s}} \Rightarrow -\frac{\zeta'_K(s)}{\zeta_K(s)} = \sum_{\alpha \in A} \lambda_\alpha \frac{\log q}{1 - \alpha^{-1} q^s} \quad (3.4)$$

Now consider the partial fraction formula :

$$\frac{1}{e^z - 1} + \frac{1}{2} = \lim_{T \rightarrow \infty} \sum_{n=-T}^T \frac{1}{z - 2\pi i n}$$

Substituting $e^z = \alpha^{-1} q^s \Rightarrow z = s \log q - \log \alpha$ we get,

$$\begin{aligned} \frac{1}{\alpha^{-1} q^s - 1} + \frac{1}{2} &= \lim_{T \rightarrow \infty} \sum_{n=-T}^T \frac{1}{s \log q - \log \alpha - 2\pi i n} \\ \frac{\log q}{\alpha^{-1} q^s - 1} + \frac{\log q}{2} &= \lim_{T \rightarrow \infty} \sum_{n=-T}^T \frac{1}{s - \frac{\log \alpha + 2\pi i n}{\log q}} \\ &= \lim_{T \rightarrow \infty} \sum_{\substack{q^\beta = \alpha \\ |\beta| \leq T}} \frac{1}{s - \beta} = \sum_{q^\beta = \alpha} \frac{1}{s - \beta} \end{aligned} \quad (3.5)$$

Putting (3.5) in equation (3.4) we get,

$$\begin{aligned} -\frac{\zeta'_K(s)}{\zeta_K(s)} &= \sum_{\alpha \in A} \lambda_\alpha \left(\frac{\log q}{2} - \sum_{q^\beta = \alpha} \frac{1}{s - \beta} \right) \\ &= \frac{\log q}{2} \sum_{\alpha \in A} \lambda_\alpha + \left(\frac{1}{s} + \frac{1}{s-1} + \sum_{\substack{\text{poles } \theta \\ \theta \neq 0,1}} \frac{1}{s - \theta} \right) - \sum_{\text{zeros}} \frac{1}{s - \rho} \\ &= \frac{1}{s} + \frac{1}{s-1} - \sum \frac{1}{s - \rho} + (g-1) \log q + \sum_{\theta \neq 0,1} \frac{1}{s - \theta} \end{aligned}$$

□

We have

$$\gamma_{K,0} = \lim_{s \rightarrow 1} \left(\frac{\zeta'_K(s)}{\zeta_K(s)} + \frac{1}{s-1} \right)$$

and using Lemma 3.1.1, we get

$$\gamma_{K,0} = \sum \frac{1}{1-\rho} - (g-1) \log q - \sum_{\theta \neq 0,1} \frac{1}{1-\theta} - 1$$

In [Ih96], Ihara deduces the following upper bound for $\gamma_{K,0}$:

Theorem 3.1.2. (Ihara) For $g > 2$ or, $g = 2$ and $q > 2$, and $\alpha_K = (g-1) \log q$ we have

$$\gamma_{K,0} \leq \left(\frac{\alpha_K + 1}{\alpha_K - 1} \right) (2 \log \alpha_K + 1 + \log q)$$

3.2 BOUNDS FOR $\gamma_{K,1}$

Differentiating (3.3) we get,

$$Z'_K(s) = -\frac{1}{s^2} - \frac{1}{(s-1)^2} + \sum \frac{1}{(s-\rho)^2} - \sum_{\theta \neq 0,1} \frac{1}{(s-\theta)^2} \quad (3.6)$$

Taking limit $s \rightarrow 1$ we get,

$$\gamma_{K,1} = 1 - \sum \frac{1}{(1-\rho)^2} + \sum_{\theta \neq 0,1} \frac{1}{(1-\theta)^2} \quad (3.7)$$

We do the same process as in the number field case, namely, we consider the integral

$$\Psi_K^{(\mu)}(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^{s-\mu}}{s-\mu} Z'_K(s) ds \quad \text{for } c \gg 0$$

for $\mu = 0$ and 1 and evaluate the expression $x\Psi_K^{(1)}(x) - \Psi_K^{(0)}(x)$ in two different ways using equation (3.6) above and another expression coming from the Euler product. Note that by $c \gg 0$, we just mean that we are considering the integral on a line $s = c$, far to the right of 1 .

Looking at the similarities of equation (3.3) to the number field case of (2.3), we see that we only need to tweak our computations a little bit, particularly, instead of the gamma factors, we need to do the computations for the sum related to the poles, rest is similar. The contribution from the

poles will be

$$\begin{aligned} & (x-1) \left[\sum_{\theta \neq 0,1} \frac{\theta(\theta-1) \log x - \theta^2 + (1-\theta)^2}{\theta^2(1-\theta)^2} \cdot \frac{x^\theta}{x-1} + \sum_{\theta \neq 0,1} \frac{1}{(1-\theta)^2} \right] \\ &= (x-1) \left[\ell(x) + \sum_{\theta \neq 0,1} \frac{1}{(1-\theta)^2} \right] \quad (\text{say}) \end{aligned}$$

Thus we will have the formula,

$$\boxed{\gamma_{K,1} = \Psi_K(x) - 1 - \frac{1+x}{1-x} \log x - \frac{1}{2}(\log x)^2 + r(x) - \ell(x)} \quad (3.8)$$

where as before,

$$\Psi_K(x) = \frac{1}{x-1} \sum_{k, N(P)^k < x} \left(\frac{x}{N(P)^k} - 1 \right) k(\log N(P))^2 \quad \text{for } x > 1 \quad (3.9)$$

and

$$r(x) = \sum_{\text{zeros}} \frac{\rho(\rho-1) \log x - \rho^2 + (1-\rho)^2}{\rho^2(1-\rho)^2} \cdot \frac{x^\rho}{x-1}$$

and

$$\ell(x) = \sum_{\substack{\text{poles} \\ \theta \neq 0,1}} \frac{\theta(\theta-1) \log x - \theta^2 + (1-\theta)^2}{\theta^2(1-\theta)^2} \cdot \frac{x^\theta}{x-1}$$

Upper bound for $\Psi_K(x)$

As in the number field case, we have

$$\begin{aligned} \Psi_K(x) &= \frac{1}{x-1} \sum_{k, N(P)^k \leq x} \left(\frac{x}{N(P)^k} - 1 \right) k(\log N(P))^2 \\ &= \frac{1}{x-1} \sum_{k, N(P)^k \leq x} \left(\frac{x}{N(P)^k} - 1 \right) (\log N(P))(\log N(P)^k) \\ &\leq (\log x) \Phi_K(x) \end{aligned} \quad (3.10)$$

Where $\Phi_K(x) = \frac{1}{x-1} \sum_{k, N(P)^k \leq x} \left(\frac{x}{N(P)^k} - 1 \right) (\log N(P))$.

This is the counterpart of our $\Psi_K(x)$ used by Ihara in [Ih96] to compute $\gamma_{K,0}$. Also note that $\Psi_K(x) \geq 0$. Ihara showed the following upper bound (e.g. see Main Lemma (see 1.5.6) and Lemma 2 of [Ih96])

$$\begin{aligned}
\Phi_K(x) &\leq \log x - \frac{\sqrt{x}-1}{\sqrt{x}+1}(\gamma_{K,0} + c_q) + \frac{2\alpha_K}{\sqrt{x}+1} + \log q \\
\Rightarrow \Psi_K(x) &\leq (\log x)^2 - \left(\frac{\sqrt{x}-1}{\sqrt{x}+1} \right) (\log x)(\gamma_{K,0} + c_q) + \frac{2\alpha_K \log x}{\sqrt{x}+1} \\
&\quad + (\log x)(\log q) \\
\Rightarrow \Psi_K(x) &\ll (\log x) \left(\log x - \gamma_{K,0} + \log q + \frac{\alpha_K}{\sqrt{x}} \right)
\end{aligned} \tag{3.11}$$

Note that, here

$$c_q = \frac{q+1}{2(q-1)} \log q$$

Upper bound for $\ell(x)$

$$\begin{aligned}
\ell(x) &= \sum_{\theta \neq 0,1} \frac{\theta(\theta-1) \log x - \theta^2 + (1-\theta)^2}{\theta^2(1-\theta)^2} \cdot \frac{x^\theta}{x-1} \\
&= \frac{\log x}{x-1} \sum_{\theta \neq 0,1} \frac{x^\theta}{\theta(\theta-1)} + \frac{1}{x-1} \sum_{\theta \neq 0,1} \left[\frac{x^\theta}{\theta^2} - \frac{x^\theta}{(1-\theta)^2} \right] \\
&\ll \log x \quad (\text{Note that } \operatorname{Re}(\theta) = 0 \text{ or } 1, \text{ and the series are abs. convg.})
\end{aligned} \tag{3.12}$$

Upper bound for $r(x)$

Since GRH holds in the Function field case, $\rho = \frac{1}{2} + i\gamma$, thus $1 - \rho = \frac{1}{2} - i\gamma = \bar{\rho}$.

We will have, like in the number field case, (and [Iha06], 1.3.11)

$$r(x) \ll \frac{\log x}{\sqrt{x}} \left(\gamma_{K,0} + \alpha_K + \frac{q+1}{2(q-1)} \log q \right) \ll \frac{\alpha_K (\log x)}{\sqrt{x}} \quad (3.13)$$

We now have all the estimates to prove our theorem :

Theorem 3.2.1. For $g > 2$ or, $g = 2$ and $q > 2$, we have

$$\gamma_{K,1} \ll (\log \alpha_K) (2 \log \alpha_K - \gamma_{K,0} + \log q + 1) \quad (3.14)$$

where $\alpha_K = (g-1) \log q$

Proof. Plugging in the bounds obtained for $\ell(x)$ in (3.12), $r(x)$ in (3.13), $\Psi_K(x)$ in (3.11) into the equation (3.8) we get :

$$\gamma_{K,1} \ll (\log x) \left(\log x - \gamma_{K,0} + \log q + \frac{\alpha_K}{\sqrt{x}} \right)$$

Choosing $x = \alpha_K^2$ we get,

$$\gamma_{K,1} \ll (\log \alpha_K) (2 \log \alpha_K - \gamma_{K,0} + \log q + 1)$$

where the last inequality follows from the bound on $\gamma_{K,0}$ due to Ihara, as in Theorem 3.1.2. \square

3.3 GENERAL CASE : BOUNDS FOR $\gamma_{K,n}$

Hence, differentiating (3.1.1) n times we get,

$$Z_K^{(n)}(s) = \frac{(-1)^n n!}{s^{n+1}} + \frac{(-1)^n n!}{(s-1)^{n+1}} + (-1)^{n+1} \sum \frac{n!}{(s-\rho)^{n+1}} + (-1)^n \sum_{\theta \neq 0,1} \frac{n!}{(s-\theta)^{n+1}} \quad (3.15)$$

and therefore, letting $\lim s \rightarrow 1$

$$\begin{aligned} -n! \gamma_{K,n} &= (-1)^n n! + (-1)^{n+1} \sum \frac{n!}{(1-\rho)^{n+1}} + (-1)^n \sum_{\theta \neq 0,1} \frac{n!}{(1-\theta)^{n+1}} \\ \Rightarrow \gamma_{K,n} &= (-1)^{n+1} + (-1)^n \sum \frac{1}{(1-\rho)^{n+1}} + (-1)^{n+1} \sum_{\theta \neq 0,1} \frac{1}{(1-\theta)^{n+1}} \end{aligned} \quad (3.16)$$

The computations are very similar. We have

$$\Psi_K(n, x) = \frac{1}{x-1} \sum_{k, N(P)^k \leq x} \left(\frac{x}{N(P)^k} - 1 \right) k^n (\log N(P))^{n+1} \quad \text{for } x > 1 \quad (3.17)$$

together with (for $n \geq 2$)

$$\Psi_K(n, x) \leq (\log x)^{n-1} \Psi_K(x) \ll (\log x)^n \left(\log x - \gamma_{K,0} + \log q + \frac{\alpha_K}{\sqrt{x}} \right) \quad (3.18)$$

The contribution from the term : $\frac{(-1)^n n!}{s^{n+1}} + \frac{(-1)^n n!}{(s-1)^{n+1}}$ can be similarly computed to be the function $f(n, x)$ as defined in Theorem 2.8.1. Also, we had $f(n, x) \ll (\log x)^{n+1}$. Similarly, since GRH is known in this case, contribution from the non-trivial zeros is

$$n!(-1)^{n+1}(x-1) \left(\sum \frac{1}{(1-\rho)^{n+1}} + r(n, x) \right)$$

where,

$$r(n, x) \ll (n!2^{n+1}) \frac{\alpha_K (\log x)^n}{\sqrt{x}}$$

The only new thing we need to compute is the contribution of the poles. Which again, looking at the similarity of the term to that of the zeros, looks like :

$$n!(-1)^n(x-1) \left(\sum \frac{1}{(1-\theta)^{n+1}} + \ell(n, x) \right)$$

where

$$\begin{aligned} \ell(n, x) &= \frac{1}{x-1} \sum_{\substack{\text{poles} \\ \theta \neq 0,1}} \left(x(-1)^n (n!) \sum_{k=0}^n \binom{n}{k} x^{\theta-1} (\log x)^{n-k} \cdot \frac{(-1)^k k!}{(\theta-1)^{k+1}} \right. \\ &\quad \left. - (-1)^n (n!) \sum_{k=0}^n \binom{n}{k} x^\theta (\log x)^{n-k} \cdot \frac{(-1)^k k!}{\theta^{k+1}} \right) \\ &\ll \frac{n! x (\log x)^n}{x-1} \sum_{k=0}^n \binom{n}{k} \left| \sum_{\substack{\text{poles} \\ \theta \neq 0,1}} \frac{1}{(\theta-1)^{k+1}} - \frac{1}{\theta^{k+1}} \right| \\ &\ll n! 2^n (\log x)^n \end{aligned}$$

Note that the first inequality follows from the fact that $\operatorname{Re}(\theta) = 0$ or 1 whereas, all the series in the second inequality is absolutely convergent.

We are now ready to generalize Theorem 3.2.1.

Theorem 3.3.1. For $g > 2$ or, $g = 2$ and $q > 2$, we have

$$\gamma_{K,n} + (-1)^n \ll \frac{2}{n!} (\log(n! 2^{n+1} \alpha_K))^n (2 \log(\alpha_K) - \gamma_{K,0} + \log q + 1 + n! 2^n)$$

As before, here $\alpha_K = (g-1) \log q$.

Proof. Putting our computations together,

$$\begin{aligned} (x-1)\Psi_K(n, x) &= f(n, x) + n!(-1)^{n+1}(x-1) \left(\sum \frac{1}{(1-\rho)^{n+1}} + r(n, x) \right) \\ &\quad + n!(-1)^n(x-1) \left(\sum \frac{1}{(1-\theta)^{n+1}} + \ell(n, x) \right) \\ \Rightarrow \Psi_K(n, x) &= \frac{f(n, x)}{(x-1)} - \left[(-1)^n \sum \frac{n!}{(1-\rho)^{n+1}} + (-1)^{n+1} \sum_{\theta \neq 0,1} \frac{n!}{(1-\theta)^{n+1}} \right] \\ &\quad + r(n, x) + \ell(n, x) \end{aligned}$$

$$\begin{aligned}
n!(\gamma_{K,n} + (-1)^n) &= -\Psi_K(n, x) + \frac{f(n, x)}{(x-1)} + r(n, x) + \ell(n, x) \\
&\ll (\log x)^n \left(\log x - \gamma_{K,0} + \log q + \frac{n!2^{n+1}\alpha_K}{\sqrt{x}} + n!2^n \right)
\end{aligned}$$

choosing $x = (n!2^{n+1}\alpha_K)^2$ to minimize the sum, we get our result

$$\gamma_{K,n} + (-1)^n \ll \frac{2}{n!} (\log(n!2^{n+1}\alpha_K))^n (2\log(\alpha_K) - \gamma_{K,0} + \log q + 1 + n!2^n)$$

□

MOMENTS OF HIGHER DERIVATIVES OF $\mathcal{L}(s, \chi)$ AT $s = 1$

4.1 PRELIMINARIES

Let K be a number field and χ be a primitive Dirichlet character on K (i.e. a primitive Hecke character with finite order). Let $L(s, \chi)$ be the L -function associated to it. In particular, when $\chi = \chi_0$, the principal character, $L(s, \chi) = \zeta_K(s)$, the Dedekind zeta function of K . The completed L -function is of the form :

$$\xi(s, \chi) = AB^{\frac{s}{2}} \Gamma\left(\frac{s+1}{2}\right)^a \Gamma\left(\frac{s}{2}\right)^{a'} \Gamma(s)^{r_2} L(s, \chi) \quad (4.1)$$

and satisfies a functional equation : $\xi(s, \chi) = \varepsilon(\chi) \xi(1-s, \bar{\chi})$, where $\varepsilon(\chi)$ is a constant of absolute value 1. A, B are constants involving $2, \pi$, the discriminant of K and the conductor f_χ . As we'll be concerned with higher derivatives, we haven't written them down explicitly, but interested reader can have a look at p.211 of [CF76] or for Hecke's original proof see [Hec83].

Also note that, here a (resp. a') is the number of real places of K where χ is ramified (resp. unramified), $r_1 = a + a'$ is the number of real places of K and r_2 is the number of complex places in K .

For $\chi \neq \chi_0$, taking the logarithmic derivative of (4.1) and using Hadamard product one can then deduce a Stark like lemma (e.g. see Lemma 2.1 of [Sta74] or p.83 of [Davoo]) :

$$\frac{L'(s, \chi)}{L(s, \chi)} = C - \frac{a}{2} \frac{\Gamma'}{\Gamma}\left(\frac{s}{2}\right) - \frac{a'}{2} \frac{\Gamma'}{\Gamma}\left(\frac{s+1}{2}\right) - r_2 \frac{\Gamma'}{\Gamma}(s) + \sum_{\rho} \left(\frac{1}{s-\rho} + \frac{1}{\rho}\right) \quad (4.2)$$

C being a constant involving log of terms in B in (4.1) etc.

For the rest of the chapter, we will denote the LHS by $\mathcal{L}(s, \chi)$, i.e.

$$\mathcal{L}(s, \chi) = \frac{L'(s, \chi)}{L(s, \chi)} \quad (\text{say})$$

On the other hand, by taking the logarithmic derivative of the Euler product of $L(s, \chi)$ we get :

$$\mathcal{L}(s, \chi) = - \sum_{P, k} \left(\frac{\chi(P)}{N(P)^s} \right)^k \log N(P) \quad (4.3)$$

Ihara, Murty and Shimura proved the following theorem in [IMS09] :

Theorem 4.1.1. (Ihara, Murty, Shimura)

If $\chi \neq \chi_0$, then

$$\mathcal{L}(1, \chi) = - \lim_{x \rightarrow \infty} \Phi_{K, \chi}(x) \quad (4.4)$$

where

$$\Phi_{K, \chi}(x) = \frac{1}{x-1} \sum_{N(P)^k \leq x} \left(\frac{x}{N(P)^k} - 1 \right) \chi(P)^k \log N(P) \quad (\text{for } x > 1)$$

Here, k is a positive integer and the sum is taken over non-archimedean primes. Under GRH, they have shown the following upper bound :

$$|\mathcal{L}(1, \chi)| < 2 \log \log \sqrt{d_\chi} + 1 - \gamma_{K,0} + O\left(\frac{\log |d_K| + \log \log d_\chi}{\log d_\chi}\right)$$

Here, $d_\chi = |d_K| N(\mathfrak{f}_\chi)$ and $\gamma_{K,0}$ is the Euler-Kronecker constant of K .

The proof of the above theorem follows its counterpart for the Dedekind zeta function, due to Ihara in [Iha06]. It is based on computing the integral

$$\Phi^{(\mu)}(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^{s-\mu}}{s-\mu} \mathcal{L}(s, \chi) ds \quad \text{for } c \gg 0$$

for $\mu = 0$ and 1 , in two different ways using the equations (4.2) and (4.3) and then estimating the terms.

Remark 4.1.2. Due to equation (4.2) and (4.3), using the same methods as in the case of $\gamma_{K,n}$ in Chapter 2, similar formulas and bounds for higher

derivatives of $\mathcal{L}(s, \chi)$ at $s = 1$ can be computed. We present some of those computations in the next sections.

4.2 AN "EXACT FORMULA" FOR $\mathcal{L}'(1, \chi)$

Differentiating equation (4.2) we get,

$$\mathcal{L}'(s, \chi) = - \sum_{\rho} \frac{1}{(s - \rho)^2} + \tilde{\Gamma}'_{\chi}(s) \tag{4.5}$$

$$\begin{aligned} \tilde{\Gamma}'_{\chi}(s) &= -\frac{a}{2} \frac{\Gamma'}{\Gamma} \left(\frac{s+1}{2} \right) - \frac{a'}{2} \frac{\Gamma'}{\Gamma} \left(\frac{s}{2} \right) - r_2 \frac{\Gamma'}{\Gamma}(s) \\ &= \frac{n}{2} \gamma + \frac{a}{2} \sum_{k=0}^{\infty} \left(\frac{2}{s+1+2k} - \frac{2}{2k+2} \right) + \frac{a'}{2} \sum_{k=0}^{\infty} \left(\frac{2}{s+2k} - \frac{2}{1+2k} \right) \\ &\quad + r_2 \sum_{k=0}^{\infty} \left(\frac{1}{s+k} - \frac{1}{1+k} \right) \end{aligned}$$

Here $\gamma = \lim_{n \rightarrow \infty} [\sum_{k=1}^n \frac{1}{k} - \ln n] \sim 0.5772\dots$ is the Euler–Mascheroni constant and $n = [K : \mathbb{Q}]$. Differentiating we get,

$$\tilde{\Gamma}'_{\chi}(s) = -a \sum_{k=0}^{\infty} \frac{1}{(s+1+2k)^2} - a' \sum_{k=0}^{\infty} \frac{1}{(s+2k)^2} - r_2 \sum_{k=0}^{\infty} \frac{1}{(s+k)^2}$$

Differentiating the Euler product in equation (4.3) we get,

$$\mathcal{L}'(s, \chi) = \sum_{P,k} k \left(\frac{\chi(P)}{N(P)^s} \right)^k (\log N(P))^2 \tag{4.6}$$

To find a similar 'exact formula' as in the case of $\gamma_{K,1}$, we evaluate the integral :

$$\Psi_{\chi}^{(\mu)}(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^{s-\mu}}{s-\mu} \mathcal{L}'(\chi, s) ds \quad \text{for } c \gg 0$$

For $\mu = 0$ and 1 in two different ways using equation (4.6) and equation (4.2) and the classical formulas:

$$\frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{y^s}{s} ds = \begin{cases} 0 & 0 < y < 1 \\ \frac{1}{2} & y = 1 \\ 1 & y > 1 \end{cases} \quad (4.7)$$

From the Euler product we get :

$$x\Psi_\chi^{(1)}(x) - \Psi_\chi^{(0)}(x) = \sum_{k, N(P)^k < x} k \left(\frac{x}{N(P)^k} - 1 \right) \chi(P)^k (\log N(P))^2 \quad (4.8)$$

Looking at the above computation, we define :

$$\Psi_\chi(x) = \frac{1}{x-1} \sum_{k, N(P)^k < x} k \left(\frac{x}{N(P)^k} - 1 \right) \chi(P)^k (\log N(P))^2 \quad \text{for } x > 1$$

(4.9)

On the other hand, from equation (4.5) we get,

$$\begin{aligned} x\Psi_\chi^{(1)}(x) - \Psi_\chi^{(0)}(x) &= \frac{x}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^{s-1}}{s-1} \left[-\sum \frac{1}{(s-\rho)^2} + \tilde{\Gamma}'_\chi(s) \right] ds \\ &\quad - \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^s}{s} \left[-\sum \frac{1}{(s-\rho)^2} + \tilde{\Gamma}'_\chi(s) \right] ds \end{aligned}$$

Computing similar contour integrals as $\gamma_{K,1}$ we see that, contribution from the \sum_ρ term is :

$$\begin{aligned} (1-x) &\left[\sum \frac{\rho(\rho-1) \log x - \rho^2 + (1-\rho)^2}{\rho^2(1-\rho)^2} \cdot \frac{x^\rho}{x-1} + \sum \frac{1}{(1-\rho)^2} \right] \\ &= (1-x) \left[r_\chi(x) + \sum \frac{1}{(1-\rho)^2} \right] \quad (\text{say}) \end{aligned} \quad (4.10)$$

For contribution from the $\tilde{\Gamma}'_\chi(s)$, we first re-write it as follows :

$$\tilde{\Gamma}'_\chi(s) = -\frac{a' + r_2}{s^2} - a \sum_{k=0}^{\infty} \frac{1}{(s+1+2k)^2} - a' \sum_{k=1}^{\infty} \frac{1}{(s+2k)^2} - r_2 \sum_{k=1}^{\infty} \frac{1}{(s+k)^2}$$

For the first term in the above equation, we have

$$\begin{aligned} & - (a' + r_2) \left[\frac{x}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^{s-1}}{s^2(s-1)} ds - \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^s}{s^3} ds \right] \\ & = - (a' + r_2) \left[\frac{x}{2\pi i} \int_{c-i\infty}^{c+i\infty} x^{s-1} \left(\frac{1}{s-1} - \frac{1}{s} - \frac{1}{s^2} \right) ds - \frac{1}{2} (\log x)^2 \right] \\ & = - (a' + r_2) \left[x - 1 - \log x - \frac{1}{2} (\log x)^2 \right] \\ & = (a' + r_2)(1-x) \left[1 - \frac{\log x}{(x-1)} - \frac{(\log x)^2}{2(x-1)} \right] \end{aligned}$$

Note that, as before, here we're using the classical formula : (for $n \geq 1$)

$$\frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{y^s}{s^{n+1}} ds = \begin{cases} 0 & 0 < y \leq 1 \\ \frac{1}{n!} (\log y)^n & y > 1 \end{cases}$$

For the rest of the terms involving series, the computations are similar to that of $\frac{1}{(s-\rho)^2}$ and we get the total contribution of $\tilde{\Gamma}'_\chi(s)$ terms to be :

$$\begin{aligned}
 & (a' + r_2)(1-x) \left[1 - \frac{\log x}{(x-1)} - \frac{(\log x)^2}{2(x-1)} \right] + \\
 & a(1-x) \left[\sum_{k=0}^{\infty} \frac{(2k+1)(2k+2) \log x - (1+2k)^2 + (2+2k)^2}{(2+2k)^2(1+2k)^2} \cdot \frac{x^{-2k-1}}{x-1} + \sum_{k=0}^{\infty} \frac{1}{(2+2k)^2} \right] + \\
 & a'(1-x) \left[\sum_{k=1}^{\infty} \frac{2k(2k+1) \log x - 4k^2 + (1+2k)^2}{4k^2(1+2k)^2} \cdot \frac{x^{-2k}}{x-1} + \sum_{k=1}^{\infty} \frac{1}{(1+2k)^2} \right] + \\
 & r_2(1-x) \left[\sum_{k=1}^{\infty} \frac{k(k+1) \log x - k^2 + (1+k)^2}{k^2(1+k)^2} \cdot \frac{x^{-k}}{x-1} + \sum_{k=1}^{\infty} \frac{1}{(1+k)^2} \right] \\
 & = (1-x) [\ell_\chi(x) - \tilde{\Gamma}'_\chi(1)] \quad (\text{say})
 \end{aligned} \tag{4.11}$$

Putting together equation (4.8), (4.10) and (4.11) we get

$$\begin{aligned}
 (x-1)\Psi_\chi(x) &= (1-x) \left[r_\chi(x) + \sum \frac{1}{(1-\rho)^2} \right] + (1-x) [\ell_\chi(x) - \tilde{\Gamma}'_\chi(1)] \\
 -\Psi_\chi(x) &= \ell_\chi(x) + r_\chi(x) - \tilde{\Gamma}'_\chi(1) + \sum \frac{1}{(1-\rho)^2} \\
 \boxed{\mathcal{L}'(1, \chi) = \Psi_\chi(x) + r_\chi(x) + \ell_\chi(x)} & \tag{4.12}
 \end{aligned}$$

Note that,

$$\begin{aligned}
 |\Psi_\chi(x)| &\leq \Psi_K(x) \\
 &\ll (\log x) \left(\log x - \gamma_{K,0} + \frac{2\alpha_K}{\sqrt{x}} \right)
 \end{aligned}$$

where the last inequality is under GRH, follows from (2.25)

Lemma 4.2.1. For $\chi \neq \chi_0$, we have (unconditionally)

$$\ell_\chi(x) = O\left(\frac{n_K(\log x)^2}{x}\right)$$

Here the implied constant is absolute.

Proof. Note that the series are absolutely convergent and thus, contribution from the series terms are : $O\left(\frac{(a+r_2)\log x}{x^2} + \frac{a'\log x}{x^3}\right) = O\left(\frac{n_K \log x}{x^2}\right)$. Whereas, the first term is $O\left(\frac{n_K(\log x)^2}{x}\right)$. \square

We are now ready to state and prove our exact formula :

Theorem 4.2.2. For $\chi \neq \chi_0$, we have, unconditionally,

$$\mathcal{L}'(\chi, 1) = \lim_{x \rightarrow \infty} \Psi_\chi(x)$$

Proof. From the above lemma, $\lim_{x \rightarrow \infty} \ell_\chi(x) = 0$. For the r_χ term, note that following the exact same steps as in the computation of $\lim_{x \rightarrow \infty} r(x)$, for $\gamma_{K,1}$ we can show that $\lim_{x \rightarrow \infty} r_\chi(x) = 0$. □

We also note that, since $r_\chi(x)$ has the same expression as that of $r(x)$ in the computation of $\gamma_{K,1}$, we get, under GRH, (writing $\rho = \frac{1}{2} + i\gamma$)

$$r_\chi(x) = \frac{\log x}{x-1} \sum \frac{x^\rho}{\rho(\rho-1)} + \frac{1}{x-1} \sum \frac{-2i\sqrt{x} \sin(\gamma \log x)}{(1-\rho)^2} \quad (4.13)$$

Thus, (for $\chi \neq \chi_0$)

$$\begin{aligned} |r(x)| &\leq \frac{\sqrt{x} \log x}{x-1} \sum \frac{1}{|\rho(\rho-1)|} + \frac{2\sqrt{x}}{x-1} \sum \frac{1}{|(1-\rho)^2|} \\ &= \frac{\sqrt{x}(\log x + 2)}{x-1} \sum \frac{1}{\rho(1-\rho)} \\ &= \frac{2\sqrt{x}(\log x + 2)}{x-1} (\mathcal{L}(\chi, 1) + \alpha_{K,\chi} + \beta_{K,\chi}) \end{aligned}$$

The last equality follows from, Theorem 2 of [IMSog]. Here,

$$\begin{cases} \alpha_{K,\chi} = \frac{1}{2} \log d_\chi & \text{where } d_\chi = |d_K| N(\mathfrak{f}_\chi) \\ \beta_{K,\chi} = -\frac{a+r_2}{2}(\gamma + \log 4\pi) - \frac{a'+r_2}{2}(\gamma + \log \pi) \end{cases}$$

\mathfrak{f}_χ being the conductor of χ and $\gamma = \gamma_{\mathbb{Q},0}$ being the Euler-Mascheroni constant. Note that, from Theorem 3 of [IMSog] we have $\mathcal{L}(\chi, 1) \ll \alpha_{K,\chi}$.

Thus we can write,

$$r_\chi(x) \ll \frac{\log x}{\sqrt{x}} (\alpha_{K,\chi}) \quad (4.14)$$

4.3 GENERALIZATION TO HIGHER DERIVATIVES

In this section we present a generalization of the limit formula as in Theorem 4.2.2. For $n \geq 1$, we look at the n -th derivative of the Euler product in (4.3) :

$$\mathcal{L}^{(n)}(s, \chi) = (-1)^{n+1} \sum_{P,k} k^n \left(\frac{\chi(P)}{N(P)^s} \right)^k (\log N(P))^{n+1} \quad (4.15)$$

Similarly differentiating (4.2) n -times,

$$\mathcal{L}^{(n)}(s, \chi) = (-1)^n n! \sum_{\rho} \frac{1}{(s - \rho)^{n+1}} + \tilde{\Gamma}_{\chi}^{(n)}(s) \quad (4.16)$$

We similarly evaluate the integral :

$$\Psi_{\chi}(\mu, n, x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{x^{s-\mu}}{s-\mu} \mathcal{L}^{(n)}(\chi, s) ds \quad \text{for } c \gg 0$$

For $\mu = 0$ and 1 in two different ways using equation (4.15) and (4.16). Thus on one hand we have,

$$\begin{aligned} & x\Psi_{\chi}(1, n, x) - \Psi_{\chi}(0, n, x) \\ &= (-1)^{n+1} \sum_{k, N(P)^k < x} k^n \left(\frac{x}{N(P)^k} - 1 \right) \chi(P)^k (\log N(P))^{n+1} \end{aligned} \quad (4.17)$$

On the other hand we can similarly compute the contribution from the \sum_{ρ} term and Γ -factor. For the non-trivial zeros we do similar contour computations. The pole at $s = 0$ (resp. $s = 1$) has residue $-\frac{n!}{\rho^{n+1}}$ (resp. $\frac{(-1)^n n!}{(1-\rho)^{n+1}}$) where as residue at $s = \rho$ (pole of order $n + 1$) is $(-1)^n \lim_{s \rightarrow \rho} \frac{d^n}{ds^n} \left(\frac{x^s}{s} \right)$ (resp. s replaced by $s - 1$ while computing $\Psi_K(1, n, x)$) so that the total contribution will be of the form $(x - 1)[r(\chi, n, x) - (-1)^n n! \sum_{\rho} \frac{1}{(1-\rho)^{n+1}}]$ where

$$r(\chi, n, x) = \frac{(-1)^n}{x - 1} \left[x \sum_{\rho} \lim_{s \rightarrow \rho} \frac{d^n}{ds^n} \left(\frac{x^{s-1}}{s-1} \right) - \sum_{\rho} \lim_{s \rightarrow \rho} \frac{d^n}{ds^n} \left(\frac{x^s}{s} \right) \right]$$

Also, following along the same lines as Theorem 2.8.1 of Chapter 2, unconditionally we have $r(\chi, n, x) \rightarrow 0$ as $x \rightarrow \infty$.

Now for the Gamma factors, first note

$$\begin{aligned} \tilde{\Gamma}_\chi^{(n)}(s) = (-1)^n n! & \left[\frac{(a' + r_2)}{s^{n+1}} + a \sum_{k=0}^{\infty} \frac{1}{(s + 1 + 2k)^{n+1}} \right. \\ & \left. + a' \sum_{k=1}^{\infty} \frac{1}{(s + 2k)^{n+1}} + r_2 \sum_{k=1}^{\infty} \frac{1}{(s + k)^{n+1}} \right] \end{aligned}$$

from the residue at $s = 0$ the $\tilde{\Gamma}_\chi^{(n)}(1)$ term will come and we will similarly be able to write, the total contribution as $(x - 1)[\ell(\chi, n, x) + \tilde{\Gamma}_\chi^{(n)}(1)]$. The main term of $\ell(\chi, n, x)$ comes from the $\frac{a'+r_2}{s^{n+1}}$ term as before and thus is $\ll \frac{(\log x)^n}{x}$, in particular $\ell(\chi, n, x) \rightarrow 0$ as $x \rightarrow \infty$. Therefore we have the following theorem.

Theorem 4.3.1. For $\chi \neq \chi_0$, we have, unconditionally

$$\mathcal{L}^{(n)}(1, \chi) = \lim_{x \rightarrow \infty} (-1)^{n+1} \Psi_K(\chi, n, x)$$

Remark 4.3.2. Note the difference in the limit formula in comparison to Theorem 2.8.1, in particular the the absence fo the $f(n, x)$ term.

Remark 4.3.3. Looking at the similarities of the computations for $\gamma_{K,m}$ for the Dedekind zeta functions of number fields, we can easily deduce some bounds for these higher coefficients $\mathcal{L}^{(n)}(1, \chi)$ as well. Here we just write them, proofs are exactly similar. Under GRH, for $|d_K| > 8$, we have

$$\mathcal{L}'(1, \chi) \ll (\log \alpha_{K,\chi})(2 \log \alpha_{K,\chi} - \gamma_{K,0})$$

whereas,

$$\mathcal{L}^{(m)}(1, \chi) \ll \frac{2^m}{m!} (A + \log \alpha_{K,\chi})^m (A + 2 \log \alpha_{K,\chi} - \gamma_{K,0})$$

Where A will be a constant, $A = O(\log(m!))$.

We will now focus our attention to the case when $K = \mathbb{Q}$ and in the following sections, study the moments of higher derivatives of $\mathcal{L}(s, \chi)$ at $s = 1$, where χ runs over all non-principal multiplicative characters of large prime conductors. Before that, let us first have a brief look at some of the rich history of the study of moments of L -functions.

4.4 MOMENTS : A BRIEF HISTORY

The distribution of values of Dirichlet L -functions $L(1, \chi)$, for variable χ has been studied extensively and has a vast literature. However the study of the same for logarithmic derivatives $L'(1, \chi)/L(1, \chi)$ is more recent. Let m be a prime and X_m denote the set of all non-principal multiplicative characters $\chi : (\mathbb{Z}/m\mathbb{Z})^\times \rightarrow \mathbb{C}^\times$ and $L(s, \chi)$ denote the corresponding Dirichlet L -function.

For any pair of non-negative integers (a, b) let $P^{(a,b)}(z) = z^a \bar{z}^b$. A result of Paley and Selberg states that (e.g. see [Pal31])

$$\frac{1}{|X_m|} \sum_{\chi \in X_m} P^{(1,1)}(L(1, \chi)) = \zeta(2) + O((\log m)^2/m)$$

This was later improved and by many authors. W. Zhang [Zha90] generalized to the case of $P^{(k,k)}$. In [IMS09], Ihara, Murty and Shimura studied the moments of the logarithmic derivative and proved the following theorem :

Theorem 4.4.1. (Ihara, Murty, Shimura)

Let m be a large prime number, and let X_m be the collection of all non-principal primitive Dirichlet characters $\chi : (\mathbb{Z}/m)^\times \rightarrow \mathbb{C}^\times$. Then

$$\frac{1}{|X_m|} \sum_{\chi \in X_m} P^{(a,b)}(L'(1, \chi)/L(1, \chi)) = (-1)^{a+b} \mu^{a,b} + O(m^{\varepsilon-1}) \quad (4.18)$$

for any $\varepsilon > 0$. In particular,

$$\lim_{m \rightarrow \infty} \frac{1}{|X_m|} \sum_{\chi \in X_m} P^{(a,b)}(L'(1, \chi)/L(1, \chi)) = (-1)^{a+b} \mu^{a,b}$$

Here $\mu^{a,b}$ is a non-negative real number defined as follows :

$$\mu^{(a,b)} = \sum_{n=1}^{\infty} \frac{\Lambda_a(n)\Lambda_b(n)}{n^2} \quad \text{where} \quad \Lambda_k(n) = \sum_{n=n_1 \cdots n_k} \Lambda(n_1) \cdots \Lambda(n_k)$$

$k > 0$ and $\Lambda(n) = \log p$, when n is a prime power and 0 otherwise (the von Mangoldt function).

In the subsequent section the author wishes to derive similar theorems on moments of the higher derivatives of $\mathcal{L}(s, \chi) = L'(s, \chi)/L(s, \chi)$ at $s = 1$. Note that, the author was not able to find a good reference that studies moments of higher derivatives of $L(s, \chi)$ at $s = 1$ but the case of $s = \frac{1}{2}$ (and fractional moments) has been studied by Conrey [CON88], Milinovich [Mil11], Heath-Brown [HB10], Soundararajan [Sou09], Sono etc. For example, Sono [Son14] recently showed :

Under GRH, for $1/2 < k < 2$ and $m \in \mathbb{Z}_{\geq 0}$ we have,

$$\frac{1}{\phi(q)} \sum_{\substack{\chi(\bmod q) \\ \chi \neq \chi_0}} P^{(k,k)} \left(L^{(m)} \left(\frac{1}{2}, \chi \right) \right) \ll (\log q)^{k^2+2km}$$

whereas for $k \geq 2$, for any $\epsilon > 0$, under GRH,

$$\frac{1}{\phi(q)} \sum_{\chi(\bmod q)}^* P^{(k,k)} \left(L^{(m)} \left(\frac{1}{2}, \chi \right) \right) \ll (\log q)^{k^2+2km+\epsilon}$$

where \sum^* is over all primitive Dirichlet characters modulo q .

However note that, methods used in the above do not seem to apply to our case. Ours is more of an extension of the work done in [IMSo9].

4.5 MOMENTS OF $\mathcal{L}'(1, \chi)$ (CONDITIONAL : UNDER GRH)

Before we dive right into our theorems, let us look at the following neat connection due to the orthogonality relations of characters.

Let $\alpha : \mathbb{N} \rightarrow \mathbb{C}$ be such that, for any $\epsilon > 0$, $\alpha(n) = O(n^\epsilon)$. Consider the Dirichlet series (absolutely convergent for $\text{Re}(s) > 1$)

$$f(s) = \sum_{n=1}^{\infty} \frac{\alpha(n)}{n^s}$$

Let $X_m^* = X_m \cup \{\chi_0\}$. For each $\chi \in X_m^*$, consider the associated series

$$f_\chi(s) = \sum_{n=1}^{\infty} \frac{\chi(n)\alpha(n)}{n^s}$$

Let $\alpha_k(n)$ denote the Dirichlet coefficient of n^{-s} in $f(s)^k$ for $k \geq 0$. Then from the orthogonality relation for characters lead to the asymptotic formula : (writing $\sigma = \text{Re}(s)$)

$$\frac{1}{|X_m^*|} \sum_{\chi \in X_m^*} P^{(a,b)}(f_\chi(s)) = \sum_{n=1}^{m-1} \frac{\alpha_a(n)\overline{\alpha_b(n)}}{n^{2\sigma}} + O_{a,b}(m^{1+\epsilon-\sigma}) \quad (4.19)$$

For any s with $\sigma > 1 + \epsilon$. In particular, taking the limit $m \rightarrow \infty$

$$\lim_{m \rightarrow \infty} \frac{1}{|X_m^*|} \sum_{\chi \in X_m^*} P^{(a,b)}(f_\chi(s)) = \sum_{n=1}^{\infty} \frac{\alpha_a(n)\overline{\alpha_b(n)}}{n^{2\sigma}}$$

Now one can ask whether this holds for $s = 1$ (or even for $\text{Re}(s) \leq 1$) when X_m^* is replaced by X_m . It turns out it depends on the analytic properties of $f(s)$ to the left of 1. For the case $f(s) = \mathcal{L}(s, \chi)$, Ihara, Murty and Shimura in [IMS09], first showed, under GRH, the error term for each χ is small. This together with bounds obtained for $\mathcal{L}(1, \chi)$ gives a formula similar to (4.18), where the main term is same and the error term is $O(\frac{(\log m)^{a+b+2}}{m})$.

To obtain the unconditional result, as stated in Theorem 4.4.1, they used Montgomery's result in [Mon71] on estimating the number of zeros in a rectangular region for $\sigma \geq 4/5$ and showed that the average value, of the absolute value of the error terms, is sufficiently small. Following in their footsteps we also first prove a conditional result.

As mentioned before, for the rest of the chapter, we are considering the case $K = \mathbb{Q}$, unless otherwise specified. Let m run over all odd prime numbers and for each m , let X_m be the collection of all non-principal

primitive Dirichlet characters $\chi : (\mathbb{Z}/m)^\times \rightarrow \mathbb{C}^\times$. As usual by $\Lambda(n)$ we will denote the von Mangoldt function :

$$\Lambda(n) = \begin{cases} \log p & \text{if } n = p^k \text{ for some prime } p \text{ and integer } k \geq 1, \\ 0 & \text{otherwise.} \end{cases}$$

Similar to [IMS09], we define

$$\Lambda_0(n) = \begin{cases} 1 & n = 1, \\ 0 & n > 1 \end{cases}$$

$$\Lambda_k(n) = \sum_{n=n_1 \cdots n_k} \Lambda(n_1) \cdots \Lambda(n_k) \quad \text{for } k > 0. \quad (4.20)$$

Note that $\Lambda_k(n) = 0$ unless the sum of exponents in the prime factorization of n is at least k . Also we have, for $1 \leq k \leq r$

$$\begin{aligned} \Lambda_k(p^r) &= \sum_{i_1 + \cdots + i_k = r} \Lambda(p^{i_1}) \cdots \Lambda(p^{i_k}) \\ &= \binom{r-1}{k-1} (\log p)^k \end{aligned} \quad (4.21)$$

Following Section 3.8 of [Iha08], we see that if n has the prime factorization $n = \prod_{i=1}^r p_i^{\alpha_i}$ then, $\Lambda_k(n)$ is the coefficient of the monomial $x_1^{\alpha_1} \cdots x_r^{\alpha_r}$ in the polynomial

$$\left(\sum_{i=1}^r (\log p_i) (x_i + x_i^2 + \cdots + x_i^{\alpha_i}) \right)^k$$

Letting $x_i = 1$ for all $i = 1, \dots, r$ we see that

$$\Lambda_k(n) \leq \left(\sum_{i=1}^r \alpha_i (\log p_i) \right)^k = (\log n)^k \quad (4.22)$$

For our purposes, we also define :

$$\ell^1 \Lambda_k(n) = \sum_{n=n_1 \cdots n_k} \Lambda(n_1) \cdots \Lambda(n_k) (\log n_1) \cdots (\log n_k) \quad \text{for } k > 0. \quad (4.23)$$

and for $k = 0$ it is equal to $\Lambda_0(n)$. Note that applying arithmetic mean is greater than or equal to geometric mean inequality we see that :

$$\prod_{i=1}^k \log n_i \leq \frac{(\log n)^k}{k^k} \quad \text{and so,}$$

$$\ell^1 \Lambda_k(n) \leq \frac{(\log n)^k}{k^k} \Lambda_k(n) \leq \frac{(\log n)^{2k}}{k^k} \quad (4.24)$$

We now have a look again on $\Psi_\chi(x)$ as in equation (4.9) or $\Psi(\chi, 1, x)$ of Theorem 4.3.1. In particular, for $K = \mathbb{Q}$, it takes the form :

$$\begin{aligned} \Psi_\chi(x) &= \frac{1}{x-1} \sum_{k, p^k < x} k \left(\frac{x}{p^k} - 1 \right) \chi(p)^k (\log p)^2 \\ &= \frac{1}{x-1} \sum_{k, p^k < x} \left(\frac{x}{p^k} - 1 \right) \chi(p^k) (\log p) (\log p^k) \\ &= \frac{1}{x-1} \sum_{n \leq x} \left(\frac{x}{n} - 1 \right) \chi(n) \Lambda(n) (\log n) \end{aligned} \quad (4.25)$$

For each pair (a, b) of non-negative integers, we define

$$\tilde{\mu}^{(a,b)} = \tilde{\mu}^{(b,a)} = \sum_{n=1}^{\infty} \frac{\ell^1 \Lambda_a(n) \ell^1 \Lambda_b(n)}{n^2} \quad (4.26)$$

Note that $\tilde{\mu}^{(0,0)} = 1$, $\tilde{\mu}^{(a,0)} = 0$ for all $a > 0$, in all other cases $\tilde{\mu} > 0$.

In particular,

$$\tilde{\mu}^{(1,1)} = \sum_{n=1}^{\infty} \left(\frac{\Lambda(n) \log(n)}{n} \right)^2$$

Theorem 4.5.1. For each pair (a, b) of non-negative integers and for $x \geq m$, we have

$$\frac{1}{|X_m|} \sum_{\chi \in X_m} P^{(a,b)}(\Psi_\chi(x)) = \tilde{\mu}^{(a,b)} + O_{a,b} \left(\frac{(\log x)^{2d}}{m} \right) \quad (4.27)$$

Here $\Psi_\chi(x)$ is as in equation(4.25) and $d = a + b + 1$.

Proof. Note that for $\chi = \chi_0$, $\Psi_{\mathbb{Q}}(x) = O((\log x)^2)$. Thus if we include the principal character in proving the theorem, it will effect the results by $O\left(\frac{(\log x)^{2a+2b}}{m}\right)$ which is less than the error term. As before, we write $X_m^* = X_m \cup \{\chi_0\}$,

$$\tilde{\mu}^{(a,b)}(x) = \frac{1}{|X_m^*|} \sum_{\chi \in X_m^*} P^{(a,b)}(\Psi_{\chi}(x)) = \frac{1}{|X_m^*|} \sum_{\chi \in X_m^*} \Psi_{\chi}(x)^a \Psi_{\bar{\chi}}(x)^b \quad (4.28)$$

For our purposes, we present in the following lemma, a general version of 4.2.2 and 4.2.3 of [IMSo9].

Lemma 4.5.2. For some $x > 1$, and $\chi \in X_m^*$ if $g_{\chi}(x) = \sum_{n \leq x} g(x, n) \chi(n)$ then,

$$\frac{1}{|X_m^*|} \sum_{\chi \in X_m^*} g_{\chi}(x)^a g_{\bar{\chi}}(x)^b = \sum_{j=1}^{m-1} \lambda^{(a)}(j, x) \lambda^{(b)}(j, x) \quad (4.29)$$

where

$$\lambda^{(k)}(j, x) = \sum_{\substack{n_1, \dots, n_k < x \\ n_1 \cdots n_k \equiv j \pmod{m}}} \prod_{i=1}^k g(x, n_i)$$

for $k \geq 1$, and for $k = 0$ define $\lambda^{(0)}(j, x) = 1$ for $j = 1$ and 0 for $j > 1$. (Recall m here is a prime number and a, b non-negative integers.)

Proof. This is a direct consequence of orthogonality relations of characters. In particular, a typical term in the sum in the LHS of (4.29) looks like

$$\left(\prod_{i=1}^a g(x, n_i) \prod_{j=1}^b g(x, m_j) \right) \chi(n_1 \cdots n_a) \bar{\chi}(m_1 \cdots m_b)$$

When summed over all χ , it has a nonzero contribution only when $(n_1 \cdots n_a) \equiv (m_1 \cdots m_b) \pmod{m}$ and hence we have our result. \square

Thus applying Lemma 4.5.2, in our case with $g(x, n) = \frac{1}{x-1} \left(\frac{x}{n} - 1\right) \Lambda(n) \log n$, we get, $g_{\chi}(x) = \Psi_{\chi}(x)$ and hence from equation (4.28)

$$\tilde{\mu}^{(a,b)}(x) = \sum_{j=1}^{m-1} \lambda^{(a)}(j, x) \lambda^{(b)}(j, x) \quad (4.30)$$

where

$$\begin{aligned}
 \lambda^{(k)}(j, x) &= \frac{1}{(x-1)^k} \sum_{\substack{n_1, \dots, n_k < x \\ n_1 \cdots n_k \equiv j \pmod{m}}} \prod_{i=1}^k \left(\frac{x}{n_i} - 1 \right) \Lambda(n_i) \log n_i \\
 &= \frac{1}{(x-1)^k} \sum_{l=0}^{\lfloor (x^k - j)/m \rfloor} \sum_{\substack{n_1, \dots, n_k < x \\ n_1 \cdots n_k = j + lm}} \prod_{i=1}^k \left(\frac{x}{n_i} - 1 \right) \Lambda(n_i) \log n_i \\
 &= \sum_{l=0}^{\lfloor (x^k - j)/m \rfloor} L^{(k)}(j + lm, x) \quad (\text{say}) \tag{4.31}
 \end{aligned}$$

here $\lfloor \cdot \rfloor$ in the upper limit of the sum, is the greatest integer function. Note that $L^{(k)}(N, x) \neq 0$ only when $N < x^k$ and in this case,

$$\begin{aligned}
 L^{(k)}(N, x) &= \frac{1}{(x-1)^k} \sum_{\substack{n_1, \dots, n_k < x \\ n_1 \cdots n_k = N}} \prod_{i=1}^k \left(\frac{x}{n_i} - 1 \right) \Lambda(n_i) \log n_i \\
 &\leq \frac{1}{N} \sum_{\substack{n_1, \dots, n_k < x \\ n_1 \cdots n_k = N}} \prod_{i=1}^k \Lambda(n_i) \log n_i \\
 &\leq \frac{1}{N} \ell^1 \Lambda_k(N) \leq \frac{(\log N)^{2k}}{k^k N} < k^k \frac{(\log x)^{2k}}{N} \tag{4.32}
 \end{aligned}$$

Thus the net contribution of the terms $l > 0$ in (4.31) is given by :

$$\begin{aligned}
 \sum_{l=1}^{\lfloor (x^k - j)/m \rfloor} L^{(k)}(j + lm, x) &< k^k \frac{(\log x)^{2k}}{m} \left(1 + \frac{1}{2} \cdots + \frac{1}{\lfloor x^k/m \rfloor} \right) \\
 &= O\left(\frac{(\log x)^{2k+1}}{m} \right)
 \end{aligned}$$

Therefore we have,

$$\lambda^{(k)}(j, x) = L^{(k)}(j, x) + O\left(\frac{(\log x)^{2k+1}}{m} \right) \tag{4.33}$$

For the main term, we also use the inequality as in (4.2.9) of [IMSo9] :

$$\text{For } x > 0 \text{ and } i, j \geq 1 \text{ we have } (x-i)(x-j) \geq (x-1)(x-ij)$$

Generalizing,

$$(x-1)^k \geq (x-n_1) \cdots (x-n_k) \geq (x-1)^{k-1} (x-n_1 \cdots n_k)$$

Thus for $n_i \geq 1$ and $n_1 \cdots n_k = j$,

$$\frac{1}{(x-1)^k} \prod_{i=1}^k \left(\frac{x}{n_i} - 1 \right) = \frac{1}{(x-1)^k} \frac{\prod_{i=1}^k (x-n_i)}{j} \leq \frac{1}{j}$$

On the other hand,

$$\begin{aligned} \frac{1}{(x-1)^k} \prod_{i=1}^k \left(\frac{x}{n_i} - 1 \right) &\geq \frac{1}{(x-1)} \frac{x-j}{j} \\ \Rightarrow 0 &\leq \frac{1}{j} - \frac{1}{(x-1)^k} \prod_{i=1}^k \left(\frac{x}{n_i} - 1 \right) \leq \frac{j-1}{j(x-1)} \end{aligned}$$

That is,

$$\frac{1}{(x-1)^k} \prod_{i=1}^k \left(\frac{x}{n_i} - 1 \right) = \frac{1}{j} + O\left(\frac{1}{x}\right) \quad (4.34)$$

Note that, in the sum of $L^{(k)}(j, x)$, since $j < m$, if we choose $x \geq m$, the condition $n_1, \dots, n_k < x$ is automatic. Thus,

$$\begin{aligned} L^{(k)}(j, x) &= \frac{1}{(x-1)^k} \sum_{n_1 \cdots n_k = j} \prod_{i=1}^k \left(\frac{x}{n_i} - 1 \right) \Lambda(n_i) \log n_i \\ &= \frac{\ell^1 \Lambda_k(j)}{j} + O\left(\frac{(\log m)^{2k}}{m}\right) \end{aligned} \quad (4.35)$$

and so,

$$\lambda^{(k)}(j, x) = \frac{\ell^1 \Lambda_k(j)}{j} + O\left(\frac{(\log x)^{2k+1}}{m}\right) \quad (4.36)$$

Plugging this in equation (4.30)

$$\tilde{\mu}^{(a,b)}(x) = \sum_{j=1}^{m-1} \frac{\ell^1 \Lambda_a(j) \ell^1 \Lambda_b(j)}{j^2} + O\left(\frac{(\log x)^{2(a+b+1)}}{m}\right) \quad (4.37)$$

Note that for $j \geq m$,

$$\begin{aligned} \sum_{j \geq m} \frac{\ell^1 \Lambda_a(j) \ell^1 \Lambda_b(j)}{j^2} &\leq \frac{1}{a^a b^b} \sum_{j \geq m} \frac{(\log j)^{2a+2b}}{j^2} \\ &= \left(\frac{(\log m)^{2a+2b}}{m}\right) \end{aligned}$$

Therefore for $x \geq m$,

$$\tilde{\mu}^{(a,b)}(x) = \tilde{\mu}^{(a,b)} + O_{a,b}\left(\frac{(\log x)^{2(a+b+1)}}{m}\right) \quad (4.38)$$

and that completes the proof. \square

We are now ready to prove the main theorem of this section, which is essentially a corollary of Theorem 4.5.1 and Theorem 4.2.2.

Theorem 4.5.3. Under GRH,

$$\frac{1}{|X_m|} \sum_{\chi \in X_m} P^{(a,b)}(\mathcal{L}'(1, \chi)) = \tilde{\mu}^{(a,b)} + O\left(\frac{(\log m)^{2(a+b+1)}}{m}\right)$$

the implicit constant depends on a, b . In particular,

$$\lim_{m \rightarrow \infty} \frac{1}{|X_m|} \sum_{\chi \in X_m} P^{(a,b)}(\mathcal{L}'(1, \chi)) = (-1)^{a+b} \tilde{\mu}^{(a,b)}$$

Proof. Note that for $K = \mathbb{Q}$, lemma 4.2 takes the form :

$$\frac{L'(s, \chi)}{L(s, \chi)} = -\frac{1}{2} \log \frac{q}{\pi} - \frac{1}{2} \frac{\Gamma'}{\Gamma} \left(\frac{s+a}{2} \right) + B(\chi) + \sum_{\rho} \left(\frac{1}{s-\rho} + \frac{1}{\rho} \right) \quad (4.39)$$

For example see p.83 of [Davoo]. Here $a = 0$ (respectively, $a = 1$) if χ is even (resp. odd) and $B(\chi) = \zeta'(0, \chi) / \zeta(0, \chi)$. The sum is over all

non-trivial zeros ρ of $L(s, \chi)$, i.e. zeros in the critical strip.

Writing $\mathcal{L}(s, \chi) = L'(s, \chi)/L(s, \chi)$ and differentiating we get

$$\mathcal{L}'(s, \chi) = - \sum_{k=0}^{\infty} \frac{1}{(s+a+2k)^2} - \sum_{\rho} \frac{1}{(s-\rho)^2} \quad (4.40)$$

From the exact formula in (4.12), Lemma 4.2.1 and equation(4.14) we get, under GRH

$$\mathcal{L}'(1, \chi) = \Psi_{\chi}(x) + O\left(\frac{\log m \log x}{\sqrt{x}} + \frac{(\log x)^2}{x}\right) \quad (4.41)$$

the implicit constant being absolute. Putting $x = m^2$ in both equation (4.41) and Theorem 4.5.1 completes the proof. \square

Remark 4.5.4. The proof Theorem 4.5.1 and the Lemma 4.5.2 suggests that we should be able to generalize these ideas for the moments of higher derivatives, $\mathcal{L}^{(n)}(1, \chi)$. We will explore more on this in a later section.

Remark 4.5.5. Note that Theorem 4.5.1 is unconditional, we are only using GRH in Theorem 4.5.3 essentially to estimate the (error) difference between $\frac{1}{|X_m|} \sum_{\chi \in X_m} P^{(a,b)}(\mathcal{L}'(1, \chi))$ and $\frac{1}{|X_m|} \sum_{\chi \in X_m} P^{(a,b)}(\Psi_{\chi}(x))$. Without GRH, it's a little more work to manage this error term, but it can be done. This is what we explore in the next section.

4.6 MOMENTS OF $\mathcal{L}'(1, \chi)$ (UNCONDITIONAL)

In this section we prove an unconditional version of Theorem 4.5.3 :

Theorem 4.6.1. For any $\epsilon > 0$, we have, unconditionally,

$$\frac{1}{|X_m|} \sum_{\chi \in X_m} P^{(a,b)}(\mathcal{L}'(1, \chi)) = \tilde{\mu}^{(a,b)} + O\left(m^{\epsilon-1}\right)$$

the implicit constant depends on a, b . In particular,

$$\lim_{m \rightarrow \infty} \frac{1}{|X_m|} \sum_{\chi \in X_m} P^{(a,b)}(\mathcal{L}'(1, \chi)) = (-1)^{a+b} \tilde{\mu}^{(a,b)}$$

Remark 4.6.2. The key difference here, is that under GRH, the individual terms for each χ in the error were small, whereas unconditionally we will show that the average of the error terms is small. In particular, we will show that for large x ,

$$\frac{1}{|X_m|} \sum_{\chi \in X_m} \left| P^{(a,b)}(\mathcal{L}'(1, \chi)) - P^{(a,b)}(\Psi_\chi(x)) \right| \ll m^{\epsilon-1}$$

To do this, much like Section 5.4 of [IMS09], we will employ zero-sum estimates of $L(s, \chi)$. Note that the above result together with Theorem 4.5.1, will give our unconditional Theorem 4.6.1.

We start with an easy inequality (This was used in 6.8 of [Iha08] and 5.3 of [IMS09]), we include a short proof as well.

Proposition 4.6.3. For any $w, z \in \mathbb{C}$ we have

$$|P^{(a,b)}(z+w) - P^{(a,b)}(z)| \leq (a+b)|w|(|z|+|w|)^{a+b-1}$$

Proof. First note that for any $n \geq 1$,

$$\begin{aligned} |(z+w)^n - z^n| &= \left| \binom{n}{1} z^{n-1}w + \cdots + \binom{n}{n} w^n \right| \\ &\leq n|w| \left(\sum_{i=1}^n \binom{n-1}{i-1} |z|^{n-i} |w|^{i-1} \right) \\ &= n|w|(|z|+|w|)^{n-1} \end{aligned}$$

where the last inequality follows from $\binom{n}{i} \leq n \binom{n-1}{i-1}$ for $1 \leq i \leq n$. Thus,

$$\begin{aligned} |P^{(a,b)}(z+w) - P^{(a,b)}(z)| &= |(z+w)^a (\overline{z+w})^b - z^a \overline{z}^b| \\ &= |(z+w)^a \overline{(z+w)}^b - z^a \overline{(z+w)}^b + z^a \overline{(z+w)}^b - z^a \overline{z}^b| \\ &\leq |z+w|^b |(z+w)^a - z^a| + |z|^a |\overline{(z+w)}^b - \overline{z}^b| \\ &\leq a|w|(|z|+|w|)^{a+b-1} + b(|z|+|w|)^a |\overline{w}|(|z|+|\overline{w}|)^{b-1} \\ &\leq (a+b)|w|(|z|+|w|)^{a+b-1} \end{aligned}$$

□

Choosing, $z = \mathcal{L}'(1, \chi)$ and $w = \Psi_\chi(x) - \mathcal{L}'(1, \chi)$ gives,

$$\begin{aligned} \left| P^{(a,b)}(\mathcal{L}'(1, \chi)) - P^{(a,b)}(\Psi_\chi(x)) \right| &\leq (a+b) |\Psi_\chi(x) - \mathcal{L}'(1, \chi)| \cdot \\ &\quad (|\Psi_\chi(x) - \mathcal{L}'(1, \chi)| + |\mathcal{L}'(1, \chi)|)^{a+b-1} \end{aligned} \quad (4.42)$$

Let us denote the unique real quadratic character in X_m by χ_1 .

We will show the following bounds :

Proposition 4.6.4.

1. For $\chi \in X_m$, $x \geq m$ and $\epsilon > 0$, we have

$$\begin{aligned} &\left| P^{(a,b)}(\mathcal{L}'(1, \chi)) - P^{(a,b)}(\Psi_\chi(x)) \right| \\ &\ll \begin{cases} (\log x)^{2a+2b-1} m^{\epsilon(a+b)} & \text{for } \chi = \chi_1 \\ ((\log x)^2 (\log m))^{(a+b-1)} |\Psi_\chi(x) - \mathcal{L}'(1, \chi)| & \text{for } \chi \neq \chi_1 \end{cases} \end{aligned}$$

2. For $x \geq m^{12}$, we have

$$\sum_{\substack{\chi \in X_m \\ \chi \neq \chi_1}} |\Psi_\chi(x) - \mathcal{L}'(1, \chi)| \ll (\log x)^{16} \quad (4.43)$$

We postpone the proof of this proposition to the end of this section as we will need several Lemmas to prove it.

Recall from the exact formula (4.12), and Lemma 4.2.1, for $\chi \in X_m$,

$$|\Psi_\chi(x) - \mathcal{L}'(1, \chi)| = |r_\chi(x)| + O\left(\frac{(\log x)^2}{x}\right) \quad (4.44)$$

where, (writing $\rho = \beta + i\gamma$)

$$\begin{aligned} |r_\chi(x)| &= \left| \sum \frac{\rho(\rho-1) \log x - \rho^2 + (1-\rho)^2}{\rho^2(1-\rho)^2} \cdot \frac{x^\rho}{x-1} \right| \\ &\leq \frac{1}{(x-1)} \sum \left(\frac{\log x}{|\rho(\rho-1)|} + \frac{1}{|\rho|^2} + \frac{1}{|(1-\rho)^2|} \right) x^\beta \end{aligned} \quad (4.45)$$

Zero-sum estimates

We now write down several lemmas to essentially estimate (4.45) and prove Prop 4.6.4. These lemmas depends on the behavior and estimates of zeros of $L(s, \chi)$ in the critical strip. To begin with, we will use the following two well-known results : (due to Gronwall, Titchmarsh, Siegel etc, e.g. see [Davoo], §14, 16 and 21)

Theorem. (A) There exists an absolute and effective positive constant c such that if $\rho = \beta + i\gamma$ is a non-trivial zero of $L(s, \chi)$ with $|\gamma| \leq T$, $T \geq 1$, then either,

$$\text{Min}(1 - \beta, \beta) > \frac{c}{\log(mT)}$$

or, $\chi = \chi_1$ and $\rho = \beta_1$ or $1 - \beta_1$ is a real simple zero satisfying $\beta_1 > \frac{1}{2}$ and $1 - \beta_1 \gg m^{-\varepsilon}$.

Theorem. (B) Let Z_χ be the set of non-trivial zeros of $L(s, \chi)$. Then

$$\#\{\beta + i\gamma \in Z_\chi : |\gamma - T| < 1\} \ll \log(m(T+2))$$

Lemma 4.6.5.

$$\sum'_{|\gamma| \leq 1} \left(\frac{\log x}{|\rho(\rho-1)|} + \frac{1}{|\rho|^2} + \frac{1}{|(1-\rho)^2|} \right) x^\beta \ll x(\log mx)(\log m)^2 \quad (4.46)$$

where \sum' is the sum over all ρ excluding the possible exceptional zero.

Proof. For $|\gamma| \leq 1$, by Theorem (A) with $T = 1$, and ρ not being exceptional, we see that $|\rho| \geq |\beta| \gg \frac{1}{\log m}$, similarly $|1 - \rho| \gg \frac{1}{\log m}$. Hence,

$$\left| \frac{1}{\rho(\rho - 1)} \right| \ll \log m, \quad \frac{1}{|\rho|^2} \ll (\log m)^2, \quad \frac{1}{|1 - \rho|^2} \ll (\log m)^2$$

Therefore,

$$\begin{aligned} & \sum'_{|\gamma| \leq 1} \left(\frac{\log x}{|\rho(\rho - 1)|} + \frac{1}{|\rho|^2} + \frac{1}{|(1 - \rho)|^2} \right) x^\beta \\ & \ll x(\log x \log m + (\log m)^2) \sum'_{|\gamma| \leq 1} 1 \\ & \ll x(\log mx)(\log m)^2 \end{aligned}$$

□

Lemma 4.6.6. For $T \geq 1$

$$\sum_{|\gamma| > T} \left(\frac{\log x}{|\rho(\rho - 1)|} + \frac{1}{|\rho|^2} + \frac{1}{|(1 - \rho)|^2} \right) x^\beta \ll \frac{x(\log x)(\log mT)}{T} \quad (4.47)$$

Proof.

$$\begin{aligned} & \sum_{|\gamma| > T} \left(\frac{\log x}{|\rho(\rho - 1)|} + \frac{1}{|\rho|^2} + \frac{1}{|(1 - \rho)|^2} \right) x^\beta \\ & \ll (x \log x) \sum_{|\gamma| > T} \frac{1}{\gamma^2} \\ & \ll (x \log x) \sum_{j=[T]}^{\infty} \frac{1}{j^2} \sum_{|\gamma - (j+1)| < 1} 1 \\ & \ll (x \log x) \sum_{j=[T]}^{\infty} \frac{\log(m(j+3))}{j^2} \\ & \ll \frac{x(\log x)(\log mT)}{T} \end{aligned}$$

□

The following result is part of the proof of a sublemma (5.4.4) of [IMS09]. We record it here as a lemma and for the sake of completion also include the proof.

Lemma 4.6.7. (Ihara, Murty and Shimura) For $T \geq 2$ and $x \geq (mT)^6$

$$\sum_{\chi \in X_m} \sum'_{\substack{\rho \in Z_\chi \\ |\gamma| \leq T}} x^\beta \ll x(\log x)^{14} \quad (4.48)$$

Proof. Let us denote

$$\tilde{S}(x, m, T) = \sum_{\chi \in X_m} \sum'_{\substack{\rho \in Z_\chi \\ |\gamma| \leq T}} x^\beta$$

The lemma is a consequence of well-known bounds for the number $N(\sigma, T, m)$ related to the number of zeros of $L(s, \chi)$ in a rectangle. In particular, for $0 \leq \sigma \leq 1$ and $T \geq 2$, define

$$\begin{cases} N(\sigma, T, \chi) = \#\{\rho = \beta + i\gamma \in Z_\chi : \beta \geq \sigma, |\gamma| \leq T\} \\ N(\sigma, T, m) = \sum_{\chi \in X_m} N(\sigma, T, \chi) \end{cases}$$

It is well known that $N(0, T, \chi) \ll T \log(mT)$ (e.g. see §16 of [Davoo]) and thus $N(0, T, m) \ll mT \log(mT)$. We will also use the following result by Montgomery (Theorem 12.1) of [Mon71], also [Mon69]:

For $\sigma \geq 4/5$ and $T \geq 2$,

$$N(\sigma, T, m) \ll (mT)^{\frac{2(1-\sigma)}{\sigma}} (\log mT)^{14} \ll (mT)^{\frac{5}{2}(1-\sigma)} (\log mT)^{14} \quad (4.49)$$

Similar result can also be found in [HJ77]. We rewrite $\tilde{S}(x, m, T)$ as

$$\tilde{S}(x, m, T) = \sum_{\chi \in X_m} \sum'_{\substack{\rho \in Z_\chi \\ |\gamma| \leq T \\ \beta < 4/5}} x^\beta + \sum_{\chi \in X_m} \sum'_{\substack{\rho \in Z_\chi \\ |\gamma| \leq T \\ 4/5 \leq \beta < 1}} x^\beta$$

The first summand is

$$\ll x^{4/5} N(0, T, m) \ll x^{4/5} (mT) (\log mT) \ll x^{4/5+1/6} \log x \ll x$$

where the last inequality is due to the imposed condition $x \geq (mT)^6$. The second summand is

$$\begin{aligned} &\leq \left| \int_{4/5}^1 x^\sigma d_\sigma N(\sigma, T, m) \right| \leq x^{4/5} N(4/5, T, m) + \left| \int_{4/5}^1 (x^\sigma \log x) N(\sigma, T, m) d\sigma \right| \\ &\ll x^{4/5} (mT)^{1/2} (\log mT)^{14} + (\log x) (mT)^{5/2} (\log mT)^{14} \int_{4/5}^1 \left(\frac{x}{(mT)^{5/2}} \right)^\sigma d\sigma \end{aligned}$$

Note that the first term is $\ll x$. Whereas the integral

$$\begin{aligned} \int_{4/5}^1 \left(\frac{x}{(mT)^{5/2}} \right)^\sigma d\sigma &= \left[\frac{\left(\frac{x}{(mT)^{5/2}} \right)^\sigma}{\log \left(\frac{x}{(mT)^{5/2}} \right)} \right]_{4/5}^1 \\ &\ll \frac{x}{(mT)^{5/2} (\log x)} \end{aligned}$$

and so the second term is $\ll x(\log mT)^{14} \ll x(\log x)^{14}$. Hence the lemma is proved. \square

Lemma 4.6.8. For $T > 1$ and $x \geq (mT)^6$ we have

$$\sum_{\chi \in X_m} \sum'_{\substack{\rho \in Z_\chi \\ |\gamma| \leq T}} \left(\frac{\log x}{|\rho(\rho-1)|} + \frac{1}{|\rho|^2} + \frac{1}{|(1-\rho)|^2} \right) x^\beta \ll x(\log x)^{16} \quad (4.50)$$

Proof. Keeping similar notation as in [IMSo9], 5.6, let us denote,

$$S(x, m, T) = \sum_{\chi \in X_m} \sum'_{\substack{\rho \in Z_\chi \\ |\gamma| \leq T}} \left(\frac{\log x}{|\rho(\rho-1)|} + \frac{1}{|\rho|^2} + \frac{1}{|(1-\rho)|^2} \right) x^\beta \quad (4.51)$$

Note that for all ρ with $\beta \leq \frac{4}{5}$, $S(x, m, T) \ll x^{4/5} (\log mx) (\log m)^2 \ll x$. This is essentially from Lemma 4.6.5 and 4.6.6. So let us focus on the zeros ρ with $\beta \geq \frac{4}{5}$. In this case, like before we divide the sum for $|\gamma| \leq 2$ and $2 < |\gamma| \leq T$. Note that, Since, $\beta \geq \frac{4}{5} > \frac{1}{3}$, we have

$$\text{Min}(\beta, 1 - \beta) = 1 - \beta > \frac{c}{\log(mT)}$$

Thus,

$$\begin{aligned} |\rho(\rho - 1)| &\geq \operatorname{Re} \rho(1 - \rho) = \beta(1 - \beta) + \gamma^2 > \frac{c_1}{\log(mT)} \\ |\rho|^2 &= \beta^2 + \gamma^2 > \frac{16}{25} \quad \text{and} \\ |1 - \rho|^2 &\geq (1 - \beta)^2 > \frac{c^2}{(\log mT)^2} \end{aligned}$$

Thus we have,

$$S(x, m, T) \ll ((\log mT)^2 + (\log mT)(\log x)) \tilde{S}(x, m, 2) + S_1(x, m, T) \quad (4.52)$$

where,

$$\begin{aligned} S_1(x, m, T) &= \sum_{\chi \in X_m} \sum'_{2 < |\gamma| \leq T} \frac{x^\beta}{\gamma^2} \\ &\leq \sum_{\substack{j \geq 0 \\ 2^{j+1} \leq T}} \frac{1}{4^j} \sum_{\chi \in X_m} \sum'_{2^j < |\gamma| \leq 2^{j+1}} x^\beta \\ &\leq \sum_{\substack{j \geq 0 \\ 2^{j+1} \leq T}} \frac{\tilde{S}(x, m, 2^{j+1})}{4^j} \ll x(\log x)^{14} \end{aligned} \quad (4.53)$$

Since, $x \geq (mT)^6$, we thus get, putting equation (4.52) and (4.53) together,

$$S(x, m, T) \ll (\log mT)(\log mTx)x(\log x)^{14} \ll x(\log x)^{16}$$

□

We are now ready to prove the proposition.

Proof of Proposition 4.6.4

1. By Lemma 4.6.5 and 4.6.6 with $T = 1$ we see that, for $\chi \neq \chi_1$,

$$r_\chi(x) \ll (\log x)(\log m)^2$$

and for $\chi = \chi_1$,

$$r_\chi(x) \ll (\log x)(\log m)^2 + (\log x)m^\epsilon \ll (\log x)m^\epsilon$$

This inequality is given by the Theorem (A), stated before Lemma 4.6.5. Putting these in (4.44) we get

$$|\Psi_\chi(x) - \mathcal{L}'(1, \chi)| \ll \begin{cases} (\log x)(\log m)^2 & \text{for } \chi \neq \chi_1 \\ (\log x)m^\epsilon & \text{for } \chi = \chi_1 \end{cases} \quad (4.54)$$

Recall that $\Psi_\chi(x) \ll (\log x)^2$, and so with the above bound for $r_\chi(x)$, we get $\mathcal{L}'(1, \chi) \ll (\log x)^2(\log m)$ for $x \geq m$ and $\chi \neq \chi_1$, whereas, $\mathcal{L}'(1, \chi_1) \ll (\log x)(\log x + m^\epsilon)$. Substituting these bounds in equation (4.42) we get, For $\chi \neq \chi_1$

$$\left| P^{(a,b)}(\mathcal{L}'(1, \chi)) - P^{(a,b)}(\Psi_\chi(x)) \right| \ll ((\log x)^2(\log m))^{(a+b-1)} |\Psi_\chi(x) - \mathcal{L}'(1, \chi)|$$

and for $\chi = \chi_1$ we get,

$$\begin{aligned} \left| P^{(a,b)}(\mathcal{L}'(1, \chi)) - P^{(a,b)}(\Psi_\chi(x)) \right| &\ll (\log x)m^\epsilon ((\log x)^2m^\epsilon)^{(a+b-1)} \\ &\ll (\log x)^{2a+2b-1}m^{\epsilon(a+b)} \end{aligned}$$

2. Putting $T = m$ in Lemma 4.6.6 we get that,

$$\sum_{\chi \in X_m} \frac{1}{x-1} \sum_{|\gamma| > m} \left(\frac{\log x}{|\rho(\rho-1)|} + \frac{1}{|\rho|^2} + \frac{1}{|(1-\rho)^2} \right) x^\beta \ll (\log x)(\log m)$$

whereas for $T = m$, Lemma 4.6.8 gives, for $x \geq m^{12}$

$$\sum_{\chi \in X_m} \frac{1}{x-1} \sum'_{|\gamma| \leq m} \left(\frac{\log x}{|\rho(\rho-1)|} + \frac{1}{|\rho|^2} + \frac{1}{|(1-\rho)^2} \right) x^\beta \ll (\log x)^{16}$$

Therefore, $\sum_{\substack{\chi \in X_m \\ \chi \neq \chi_1}} |\Psi_\chi(x) - \mathcal{L}'(1, \chi)| \ll (\log x)^{16}$.

□

Proof of Theorem 4.6.1

Putting $x = m^{12}$ in Proposition 4.6.4, we get

$$\begin{aligned} & \sum_{\chi \in X_m} \left| P^{(a,b)}(\mathcal{L}'(1, \chi)) - P^{(a,b)}(\Psi_\chi(x)) \right| \\ & \ll (\log m)^{2a+2b-1} m^{\epsilon(a+b)} + (\log m)^{3(a+b-1)} \sum_{\substack{\chi \in X_m \\ \chi \neq \chi_1}} |\Psi_\chi(m^{12}) - \mathcal{L}'(1 - \chi)| \\ & \ll (\log m)^{2a+2b-1} m^{\epsilon(a+b)} + (\log m)^{3(a+b-1)+16} \ll m^{\epsilon'} \end{aligned}$$

Hence we have,

$$\begin{aligned} \frac{1}{|X_m|} \sum_{\chi \in X_m} P^{(a,b)}(\mathcal{L}'(1, \chi)) &= \frac{1}{|X_m|} \sum_{\chi \in X_m} P^{(a,b)}(\Psi_\chi(m^{12})) + O(m^{\epsilon'-1}) \\ &= \tilde{\mu}^{(a,b)} + O(m^{\epsilon'-1}) \end{aligned}$$

Note that the last equality follows from Theorem 4.5.1 with $x = m^{12}$. \square

4.7 MOMENTS OF HIGHER DERIVATIVES $\mathcal{L}^{(n)}(1, \chi)$

We will now generalize the results in section 4.5 to higher derivatives. For this we look back at Theorem 4.3.1. Recall we defined,

$$\Psi_K(\chi, r, x) = \frac{1}{x-1} \sum_{k, N(P)^k < x} k^r \left(\frac{x}{N(P)^k} - 1 \right) \chi(P)^k (\log N(P))^{r+1}$$

In particular, for $K = \mathbb{Q}$, it takes the form

$$\begin{aligned} \Psi(\chi, r, x) &= \Psi_{\mathbb{Q}}(\chi, r, x) \\ &= \frac{1}{x-1} \sum_{k, p^k < x} k^r \left(\frac{x}{p^k} - 1 \right) \chi(p)^k (\log p)^{r+1} \\ &= \frac{1}{x-1} \sum_{k, p^k < x} \left(\frac{x}{p^k} - 1 \right) \chi(p^k) (\log p) (\log p^k)^r \\ &= \frac{1}{x-1} \sum_{n < x} \left(\frac{x}{n} - 1 \right) \chi(n) \Lambda(n) (\log n)^r \end{aligned}$$

Therefore we define, for $k > 0, r \geq 0$

$$\ell^r \Lambda_k(n) = \sum_{n_1 n_2 \dots n_k = n} \left(\prod_{i=1}^k \Lambda(n_i) (\log n_i)^r \right) \quad (4.55)$$

whereas, for $k = 0, \ell^r \Lambda_0(n) = \Lambda_0(n)$. With this, define, for $r \geq 0$,

$$\mu^{(a,b)}(r) = \sum_{j=1}^{\infty} \frac{\ell^r \Lambda_a(j) \ell^r \Lambda_b(j)}{j^2} \quad (4.56)$$

In particular, $\mu^{(a,b)}(0) = \mu^{(a,b)}$ as in 4.1.5 of [IMS09] or Theorem 4.4.1, whereas $\mu^{(a,b)}(1) = \tilde{\mu}^{(a,b)}$ as defined in equation (4.26) in the previous sections. We are now ready to state a generalization of Theorem 4.5.1.

Theorem 4.7.1. For each pair (a, b) of non-negative integers, $r \geq 0$ and for $x \geq m$, we have

$$\frac{1}{|X_m|} \sum_{\chi \in X_m} P^{(a,b)}(\Psi(\chi, r, x)) = \mu^{(a,b)}(r) + O_{a,b} \left(\frac{(\log x)^{(r+1)d+2}}{m} \right) \quad (4.57)$$

Here $d = a + b$.

Proof. The proof follows the $r = 1$ case in Theorem 4.5.1 very closely. Applying Lemma 4.5.2 with $g(x, n) = \frac{1}{(x-1)} \left(\frac{x}{n} - 1 \right) \Lambda(n) (\log n)^r$ we get,

$$\frac{1}{|X_m^*|} \sum_{\chi \in X_m^*} P^{(a,b)}(\Psi(\chi, r, x)) = \sum_{j=1}^{m-1} \lambda^{(a)}(j, x) \lambda^{(b)}(j, x) \quad (4.58)$$

where

$$\begin{aligned} \lambda^{(k)}(j, x) &= \sum_{\substack{n_1, \dots, n_k < x \\ n_1 \cdots n_k \equiv j \pmod{m}}} \prod_{i=1}^k \frac{1}{(x-1)} \left(\frac{x}{n_i} - 1 \right) \Lambda(n_i) (\log n_i)^r \\ &= \frac{1}{(x-1)^k} \sum_{l=0}^{\lfloor (x^k-j)/m \rfloor} \sum_{\substack{n_1, \dots, n_k < x \\ n_1 \cdots n_k = j+lm}} \prod_{i=1}^k \left(\frac{x}{n_i} - 1 \right) \Lambda(n_i) (\log n_i)^r \end{aligned} \quad (4.59)$$

As before we write it as

$$\lambda^{(k)}(j, x) = \sum_{l=0}^{\lfloor (x^k-j)/m \rfloor} L^{(k)}(j+lm, x)$$

and show that the total contribution from $l > 0$ terms is small. For this we note that, $L^{(k)}(N, x) \neq 0$ only when $N < x^k$ and in this case,

$$\begin{aligned} L^{(k)}(N, x) &\leq \frac{1}{N} \sum_{\substack{n_1, \dots, n_k < x \\ n_1 \cdots n_k = N}} \left(\prod_{i=1}^k \Lambda(n_i) (\log n_i)^r \right) \\ &\leq \frac{1}{N} \ell^r \Lambda_k(N) \leq k^k \frac{(\log x)^{(r+1)k}}{N} \end{aligned}$$

where the last inequality follows from : (similar to (4.24))

$$\ell^r \Lambda_k(n) \leq \frac{(\log n)^{rk}}{k^{rk}} \Lambda_k(n) \leq \frac{(\log n)^{(r+1)k}}{k^{rk}}$$

and therefore,

$$\begin{aligned} \sum_{l=1}^{\lfloor (x^k-j)/m \rfloor} L^{(k)}(j+lm, x) &< k^k \frac{(\log x)^{(r+1)k}}{m} \left(1 + \frac{1}{2} + \cdots + \frac{1}{\lfloor x^k/m \rfloor} \right) \\ &= O\left(\frac{(\log x)^{(r+1)k+1}}{m} \right) \end{aligned}$$

For the $l = 0$ term using (4.34) and for $x \geq m$ we have

$$L^{(k)}(j, x) = \frac{\ell^r \Lambda_k(j)}{j} + O\left(\frac{(\log m)^{(r+1)k}}{m} \right) \quad (4.60)$$

and hence,

$$\lambda^{(k)}(j, x) = \frac{\ell^r \Lambda_k(j)}{j} + O\left(\frac{(\log x)^{(r+1)k+1}}{m}\right)$$

and so, (writing $d = a + b$)

$$\frac{1}{|X_m^*|} \sum_{\chi \in X_m^*} P^{(a,b)}(\Psi(\chi, r, x)) = \sum_{j=1}^{m-1} \frac{\ell^r \Lambda_a(j) \ell^r \Lambda_b(j)}{j^2} + O\left(\frac{(\log x)^{(r+1)d+2}}{m}\right)$$

which, together with the inequality :

$$\begin{aligned} \sum_{j \geq m} \frac{\ell^r \Lambda_a(j) \ell^r \Lambda_b(j)}{j^2} &\leq \frac{1}{(a^a b^b)^r} \sum_{j \geq m} \frac{(\log j)^{(r+1)(a+b)}}{j^2} \\ &= O\left(\frac{(\log m)^{(r+1)(a+b)}}{m}\right) \end{aligned}$$

proves the theorem. □

Now applying Theorem 4.3.1 directly, with $x = m^2$ gives,

$$\begin{aligned} &\frac{1}{|X_m|} \sum_{\chi \in X_m} P^{(a,b)}(\mathcal{L}^{(r)}(1, \chi)) \\ &= \frac{1}{|X_m|} \sum_{\chi \in X_m} P^{(a,b)}\left((-1)^{r+1} \Psi(\chi, r, m^2)\right) + O\left(\frac{(\log m)^{rd}}{m^d}\right) \\ &= (-1)^{(r+1)d} \frac{1}{|X_m|} \sum_{\chi \in X_m} P^{(a,b)}(\Psi(\chi, r, m^2)) + O\left(\frac{(\log m)^{rd}}{m^d}\right) \\ &= (-1)^{(r+1)d} \mu^{(a,b)}(r) + O\left(\frac{(\log m)^{(r+1)d+2}}{m}\right) \end{aligned}$$

Here the last line follows from Theorem 4.7.1 with $x = m^2$. Therefore we have the following general version of Theorem 4.5.3 :

Theorem 4.7.2. Under GRH,

$$\frac{1}{|X_m|} \sum_{\chi \in X_m} P^{(a,b)}(\mathcal{L}^{(r)}(1, \chi)) = (-1)^{(r+1)d} \mu^{(a,b)}(r) + O\left(\frac{(\log m)^{(r+1)d+2}}{m}\right)$$

Here $d = a + b$ and the implicit constant depends on a, b . In particular,

$$\lim_{m \rightarrow \infty} \frac{1}{|X_m|} \sum_{\chi \in X_m} P^{(a,b)}(\mathcal{L}'(1, \chi)) = ((-1)^{(r+1)d} \mu^{(a,b)}(r))$$

Remark 4.7.3. Note that for an unconditional version of the above theorem, we need to have a closer look at the general $r_\chi(n, x)$ term. We have,

$$\begin{aligned} r_\chi(n, x) &= \frac{(-1)^n n!}{x-1} \left[x \sum_{\rho} \lim_{s \rightarrow \rho} \frac{d^n}{ds^n} \left(\frac{x^{s-1}}{s-1} \right) - \sum_{\rho} \lim_{s \rightarrow \rho} \frac{d^n}{ds^n} \left(\frac{x^s}{s} \right) \right] \\ &= \frac{(-1)^n n!}{x-1} \sum_{\rho} \left[x \sum_{k=0}^n \binom{n}{k} x^{\rho-1} (\log x)^{n-k} \cdot \frac{(-1)^k k!}{(\rho-1)^{k+1}} \right. \\ &\quad \left. - \sum_{k=0}^n \binom{n}{k} x^{\rho} (\log x)^{n-k} \cdot \frac{(-1)^k k!}{\rho^{k+1}} \right] \\ &\ll \frac{(\log x)^n}{x} \sum_{\rho} \left| \frac{x^{\rho}}{\rho(\rho-1)} \right| \quad (\text{The implicit constant depends on } n.) \end{aligned}$$

This reduces the case to that shown by Ihara, Murty and Shimura, see Sublemma 5.4.4 of [IMSog] with the difference that the implicit constant depends on n as well, apart from a and b . Therefore we have

Theorem 4.7.4. For any $\epsilon > 0$, we have, unconditionally,

$$\frac{1}{|X_m|} \sum_{\chi \in X_m} P^{(a,b)}(\mathcal{L}^{(r)}(1, \chi)) = (-1)^{(r+1)d} \mu^{(a,b)}(r) + O(m^{\epsilon-1})$$

The implicit constant depends on a, b and r . In particular,

$$\lim_{m \rightarrow \infty} \frac{1}{|X_m|} \sum_{\chi \in X_m} P^{(a,b)}(\mathcal{L}'(1, \chi)) = ((-1)^{(r+1)d} \mu^{(a,b)}(r))$$

DISTRIBUTION

5.1 PRELIMINARIES

For most of this chapter K is either \mathbb{Q} or an imaginary quadratic number field. In particular K has exactly one Archimedean prime denoted by \wp_∞ . Let χ run over all Dirichlet characters on K whose conductor (the non-archimedean part) is a prime divisor, such that $\chi(\wp_\infty) = 1$.

The average of a complex valued function $\phi(\chi)$, over a family of χ as defined above, is taken as follows :

$$\text{Avg}_\chi \phi(\chi) = \lim_{m \rightarrow \infty} \text{Avg}_{N(\mathfrak{f}) \leq m} \phi(\chi)$$

where

$$\text{Avg}_{N(\mathfrak{f}) \leq m} \phi(\chi) = \frac{\sum_{N(\mathfrak{f}) \leq m} \left(\sum_{\chi \equiv \mathfrak{f}} \phi(\chi) \right) / \sum_{\chi \equiv \mathfrak{f}} 1}{\sum_{N(\mathfrak{f}) \leq m} 1}$$

For the above setting, the following distribution theorem was proved by Ihara in [Ihara8] :

Theorem 5.1.1. (Ihara) For K as above and for $\sigma = \text{Re}(s) > 1$, there exists a real valued function $M_\sigma : \mathbb{C} \rightarrow \mathbb{R}$ satisfying, $M_\sigma(w) \geq 0$, is C^∞ in w and $\int_{\mathbb{C}} M_\sigma(w) |dw| = 1$, such that

$$\text{Avg}_\chi \Phi \left(\frac{L'(\chi, s)}{L(\chi, s)} \right) = \int_{\mathbb{C}} M_\sigma(w) \Phi(w) |dw| \quad (5.1)$$

holds for any continuous function Φ of \mathbb{C} . Moreover,

$$\text{Avg}_\chi \psi_z \left(\frac{L'(\chi, s)}{L(\chi, s)} \right) = \tilde{M}_\sigma(z)$$

where $\tilde{M}_\sigma(z)$ comes from the Fourier transform of $M_\sigma(z)$ in the sense that

$$\tilde{M}_\sigma(z) = \int_{\mathbb{C}} M_\sigma(w) \psi_z(w) |dw|$$

here $\psi_z : \mathbb{C} \rightarrow \mathbb{C}^1$ is the additive character $\psi_z(w) = \exp(i \cdot \operatorname{Re}(\bar{z}w))$

Remark 5.1.2. Note that Ihara shows this more generally, in the sense that he considers certain function fields of one variable over a finite field (the theorem is true in this case for $\sigma > 3/4$), $K = \mathbb{Q}$ and χ runs over characters of the form $N(\wp)^{-\tau i}$ and when K is a number field having more than one archimedean prime and χ runs over all “normalized unramified Grössencharacters” of K modifying the definition of average accordingly.

Remark 5.1.3. In a later paper [IM11], Ihara together with Matsumoto showed the above theorem for $\sigma \geq \frac{1}{2} + \epsilon$, under GRH and with “mild” conditions on the test function namely, $\Phi(w) \ll e^{a|w|}$ holds for some $a > 0$, or Φ is the characteristic function of either a compact subset or the complement of such a subset.

Similar distribution result for real characters was also proved by Mourtada, my academic sister, in her thesis, see [Mou13], [MM15]. She showed, for a fundamental discriminant D and a real character χ_D attached to D , let

$$N(y) := \{|D| \leq y : D \text{ is a fundamental discriminant}\}$$

then the following theorem holds.

Theorem 5.1.4. (Mourtada, Murty) Let $\sigma > \frac{1}{2}$ and assume GRH. Then there exists a density function $\mathcal{Q}_\sigma(x)$ such that

$$\lim_{y \rightarrow \infty} \frac{1}{N(y)} \sum_{\substack{|D| \leq y \\ D \text{ fund. disc.}}} \Phi\left(\frac{L'(\sigma, \chi_D)}{L(\sigma, \chi_D)}\right) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \mathcal{Q}_\sigma(x) \Phi(x) dx$$

holds for any bounded continuous function Φ on \mathbb{R} . It also holds when Φ is the characteristic function of either a compact subset of \mathbb{R} or the complement of such a subset.

Our goal in this chapter is to deduce similar distribution theorems for higher derivatives of the logarithmic derivative of Dirichlet L -functions. We have been able to prove results similar to that of Theorem 5.1.1, for

$\sigma > 1$. Whereas generalization of the later developments are still a work in progress. We'll show in a concluding section where these later developed techniques fail if we consider higher derivatives.

5.2 DISTRIBUTION FUNCTIONS : SOME BACKGROUND

In this section we present some background related to distribution. The results presented are based on the paper [JW35] of Jessen and Wintner.

Let \mathbb{R}^k be a k -dimensional Euclidean space and $\mathbf{x} = (x_1, \dots, x_k)$ be a variable point.

Definition 5.2.1. A completely additive, non-negative set function $\phi(E)$ defined for all Borel sets E in \mathbb{R}^k and having the value 1 for $E = \mathbb{R}^k$ will be called a *distribution function* in \mathbb{R}^k .

Notation. An integral with respect to ϕ will be denoted by

$$\int_E f(\mathbf{x})\phi(d\mathbf{x})$$

and is to be understood in the Lebesgue-Radon (or Lebesgue-Stieltjes) sense.

Definition 5.2.2. A set E is called a *continuity set* of ϕ if $\phi(E^\circ) = \phi(\bar{E})$ where E° denotes the set formed by all interior points of E and \bar{E} is the closure of E .

Definition 5.2.3. A sequence of distribution functions ϕ_n is said to be *convergent* if there exists a distribution function ϕ such that $\phi_n(E) \rightarrow \phi(E)$ for all continuity sets E of the limit function ϕ , which is then unique. We will use the notation $\phi_n \rightarrow \phi$.

Proposition 5.2.4. A sequence of distribution functions $\{\phi_n\}$ converges to a distribution function ϕ if and only if

$$\int_{\mathbb{R}^k} f(\mathbf{x})\phi_n(d\mathbf{x}) \rightarrow \int_{\mathbb{R}^k} f(\mathbf{x})\phi(d\mathbf{x})$$

holds for all continuous and bounded functions f . Moreover, if $\phi_n \rightarrow \phi$ then,

$$\int_{\mathbb{R}^k} f(\mathbf{x})\phi(d\mathbf{x}) \leq \liminf \int_{\mathbb{R}^k} f(\mathbf{x})\phi_n(d\mathbf{x})$$

holds for every non-negative, continuous function f .

Definition 5.2.5. If ϕ_1 and ϕ_2 are two distribution functions, then we define a new distribution function as their convolution, as follows :

$$\phi_1 * \phi_2(E) := \int_{\mathbb{R}^k} \phi_1(E - \mathbf{x})\phi_2(d\mathbf{x})$$

for every Borel set E . Here $E - \mathbf{x}$ denotes the set obtained from E by the translation $-\mathbf{x}$.

Note that one can show, $\phi_1 * \phi_2 = \phi_2 * \phi_1$

Definition 5.2.6. The spectrum $S = S(\phi)$ of a distribution function ϕ is the set of points $\mathbf{x} \in \mathbb{R}^k$ for which $\phi(E) > 0$ for any set E containing \mathbf{x} as an interior point. We note that S is always a non-empty closed set.

Definition 5.2.7. The point spectrum $P = P(\phi)$ is the set of points x such that $\phi(\{x\}) > 0$.

Definition 5.2.8. A distribution function is called continuous if $P(\phi)$ is empty, and is called absolutely continuous if $\phi(E) = 0$ for all Borel sets E of measure 0.

Proposition 5.2.9. [JW35] A distribution function ϕ is absolutely continuous iff there exists a Lebesgue integrable point function $D(x)$ in \mathbb{R} such that

$$\phi(E) = \int_E D(x)dx$$

for any Borel set E . We call $D(x)$ the density function of ϕ .

5.3 CONSTRUCTION OF $M_{\sigma,P}$ FUNCTIONS

Let P be any finite set of non-archimedean primes of K and set

$T_P := \prod_{\wp \in P} \mathbb{C}^1$, where \mathbb{C}^1 denotes, $\{z : |z| = 1\}$.

The following lemma was proved in [Iha08], (Lemma 4.3.1)

Lemma 5.3.1. Let K be as above and χ run over a family as above, excluding those characters such that $\mathfrak{f}_\chi \in P$. For each such χ , let $\chi_P = (\chi(\wp))_{\wp \in P} \in T_P$. Then $(\chi_P)_\chi$ is uniformly distributed on T_P . i.e. for any continuous function $\Psi : T_P \rightarrow \mathbb{C}$ we have,

$$\text{Avg}_\chi \Psi(\chi_P) = \int_{T_P} \Psi(t_P) d^*t_P$$

where $d^*t_\wp = (2\pi i t_\wp)^{-1} dt_\wp$ is the normalized Haar measure on the t_\wp -unit circle.

Remark 5.3.2. The above lemma is the key ingredient of our results. The idea is to make suitable change of variables in the above lemma, so that from the Jacobian a density function can be extracted.

Now define, $g_{\sigma,P} : T_P \rightarrow \mathbb{C}$ by

$$g_{\sigma,P}(t_P) = \sum_{\wp \in P} g_{\sigma,\wp}(t_\wp) \quad \text{where} \quad g_{\sigma,\wp}(t_\wp) = \frac{t_\wp N_\wp^\sigma (\log N_\wp)^2}{(t_\wp - N_\wp^\sigma)^2}$$

where $t_P = (t_\wp)_{\wp \in P}$, in particular, $|t_\wp| = 1$.

For any character χ of K which is unramified over P , let

$$L_P(\chi, s) = \prod_{\wp \in P} (1 - \chi(\wp) N_\wp^{-s})^{-1}$$

Then we have,

$$\mathcal{L}_P(\chi, s) := \frac{L'_P(\chi, s)}{L_P(\chi, s)} = \sum_{\wp \in P} -\frac{\chi(\wp) N_\wp^{-s} \log N_\wp}{(1 - \chi(\wp) N_\wp^{-s})}$$

and so,

$$\mathcal{L}'_P(\chi, s) = \frac{d}{ds} \frac{L'_P(\chi, s)}{L_P(\chi, s)} = \sum_{\wp \in P} \frac{\chi(\wp) N_{\wp}^{-s} (\log N_{\wp})^2}{(1 - \chi(\wp) N_{\wp}^{-s})^2} = g_{\sigma,P}(\chi_P N P^{-it})$$

where $t = \text{Im}(s)$ and $\chi_P N P^{-it} = (\chi(\wp) N_{\wp}^{-it})_{\wp \in P}$.

For $(\mathbf{f}_{\chi}, P) = 1$, since, $\{\chi_P\}_{\chi}$ is uniformly distributed on T_P , so is its translate $\{\chi_P N P^{-it}\}_{\chi}$. Thus for any continuous function Φ on \mathbb{C} , by the above lemma 5.3.1, applied to $\Psi = \Phi \circ g_{\sigma,P}$, we get

$$\text{Avg}_{\chi} (\Phi (\mathcal{L}'_P(\chi, s))) = \int_{T_P} \Phi(g_{\sigma,P}(t_P)) d^* t_P \quad (5.2)$$

We first note the following.

Lemma 5.3.3. For fixed s , with $\sigma = \text{Re}(s) > 1$, and for $P = P_y = \{\wp : N_{\wp} \leq y\}$, as $y \rightarrow \infty$, $\mathcal{L}'_P(\chi, s)$ tends uniformly to $\mathcal{L}'(\chi, s)$.

Proof. For any χ we have,

$$|\mathcal{L}'(\chi, s) - \mathcal{L}'_P(\chi, s)| \leq \sum_{\wp \notin P} \frac{N_{\wp}^{\sigma} \log N_{\wp}^2}{(N_{\wp}^{\sigma} - 1)^2}$$

Thus letting $y \rightarrow \infty$, RHS tends to 0. \square

Theorem 5.3.4. Let $\sigma > 0$. Then there exists a function $M_{\sigma,P} : \mathbb{C} \rightarrow \mathbb{R}$ such that, for any continuous function $\Phi(w)$ on \mathbb{C} ,

$$\int_{\mathbb{C}} M_{\sigma,P}(w) \Phi(w) |dw| = \int_{T_P} \Phi(g_{\sigma,P}(t_P)) d^* t_P$$

where $w = x + iy$ and $|dw| = (2\pi)^{-1} dx dy$, and $d^* t_P$ is the normalized Haar measure on T_P . This $M_{\sigma,P}$ function is compactly supported and satisfies the following properties :

1. $M_{\sigma,P}(w) \geq 0$,
2. $M_{\sigma,P}(\bar{w}) = M_{\sigma,P}(w)$,

$$3. \int_{\mathbb{C}} M_{\sigma,P}(w) |dw| = 1.$$

Proof. We first consider the case when $|P| = 1$, say $P = \{\varphi\}$. Let $T_{\varphi} = \mathbb{C}^1$ and write $t_{\varphi} = e^{i\theta}$ and so $d^*t_{\varphi} = \frac{1}{2\pi}d\theta$.

We consider the open unit disc, $z = re^{i\theta}$ for $0 \leq r < 1$ and $0 \leq \theta < 2\pi$. Consider the map

$$w = w(z) = \frac{(\log N_{\varphi})^2 re^{i\theta}}{(1 - re^{i\theta})^2} = \frac{A re^{i\theta}}{(1 - re^{i\theta})^2}$$

For computational brevity we'll write $A = (\log N_{\varphi})^2$ as this is just a constant. Let ρ be a real number such that, $N_{\varphi}^{-\sigma} < \rho < 1$ and let B_{ρ} be the region surrounded by the curve :

$$w = \frac{A \rho e^{i\theta}}{(1 - \rho e^{i\theta})^2}$$

Thus $w = w(z)$ gives a one-to-one correspondence between the region $B_{\sigma,\varphi}$ and the disc $C_{\rho} := \{z : |z| < \rho\}$.

Let us now compute the Jacobian of this mapping. We see that,

$$w(z) = A \frac{r \cos \theta - 2r^2 + r^3 \cos \theta}{|1 - re^{i\theta}|^4} + i A \frac{r \sin \theta - r^3 \sin \theta}{|1 - re^{i\theta}|^4} = U + iV \quad (\text{say}) \quad (5.3)$$

Thus the Jacobian is given by :

$$J = \begin{vmatrix} \frac{\partial U}{\partial r} & \frac{\partial U}{\partial \theta} \\ \frac{\partial V}{\partial r} & \frac{\partial V}{\partial \theta} \end{vmatrix} = \frac{A^2 r |1 + re^{i\theta}|^2}{|1 - re^{i\theta}|^6}$$

Note : This Jacobian was computed using a computer algebra system.

And so we have

$$\begin{aligned} \int_{T_\varphi} \Phi(g_{\sigma,\varphi}(t_\varphi)) d^*t_\varphi &= \frac{1}{2\pi} \int_0^{2\pi} \Phi\left(\frac{e^{i\theta} N_\varphi^\sigma (\log N_\varphi)^2}{(e^{i\theta} - N_\varphi^\sigma)^2}\right) d\theta \\ &= \frac{1}{2\pi} \int_0^{2\pi} \Phi\left(\frac{e^{i\theta} N_\varphi^{-\sigma} (\log N_\varphi)^2}{(1 - N_\varphi^{-\sigma} e^{i\theta})^2}\right) d\theta \\ &= \frac{1}{2\pi} \int \int_{B_{\sigma,\varphi}} \Phi(w) \delta(r - N_\varphi^{-\sigma}) J^{-1} dU dV \end{aligned}$$

where $\delta(\cdot)$ denotes the Dirac delta distribution and $w = U + iV$. Therefore we define $M_{\sigma,\varphi}(w)$ in the following way :

$$M_{\sigma,\varphi}(w) = J^{-1} \delta(r - N_\varphi^{-\sigma}) = \frac{|1 - re^{i\theta}|^6}{(\log N_\varphi)^2 |1 + re^{i\theta}|^2} \frac{\delta(r - N_\varphi^{-\sigma})}{r} \quad (5.4)$$

for $w \in B_{\sigma,\varphi}$ and $M_{\sigma,\varphi}(w) = 0$ otherwise. Plugging this in, we get

$$\int_{T_\varphi} \Phi(g_{\sigma,\varphi}(t_\varphi)) d^*t_\varphi = \int_{\mathbb{C}} M_{\sigma,\varphi}(w) \Phi(w) |dw|$$

This proves the case $P = \{\varphi\}$. For the general case, we define the function using convolution product. That is,

$$M_{\sigma,P}(w) = *_{\varphi \in P} M_{\sigma,\varphi}(w)$$

in other words, for $P = P' \cup \{\varphi\}$ define

$$M_{\sigma,P}(w) = \int_{\mathbb{C}} M_{\sigma,P'}(w') M_{\sigma,\varphi}(w - w') |dw'| \quad (5.5)$$

Note that, for any open set $U \subseteq \mathbb{C}$ we get,

$$\int_U M_{\sigma,P}(w) |dw| = \text{Vol}(g_{\sigma,P}^{-1}(U)) \quad (5.6)$$

where the volume is with respect to d^*t_P and thus $\int_{\mathbb{C}} M_{\sigma,P}(w) |dw| = 1$. The Haar measure is normalized, i.e the total volume of T_P is 1. \square

Our next Goal is to show, for $P = P_y = \{\varphi : N_\varphi \leq y\}$ as before, as $y \rightarrow \infty$, $M_{\sigma,P_y}(w)$ converges to a function $M_\sigma(w)$ uniformly in w .

Proposition 5.3.5. If $P = P_y$ and $y \rightarrow \infty$, for $\sigma > 1/2$, $M_{\sigma,P}(w)$ converges to $M_\sigma(w)$ uniformly in w . The limit, $M_\sigma(w)$ is therefore continuous in w and non-negative.

Proof. For $\wp \notin P$ we have, (writing $N\wp^{-\sigma} = q$)

$$\begin{aligned} |M_{\sigma,P \cup \{\wp\}}(w) - M_{\sigma,P}(w)| &= \left| \frac{1}{2\pi} \frac{1}{(\log N\wp)^2} \int_0^{2\pi} \frac{|1 - qe^{i\theta}|^6}{|1 + qe^{i\theta}|^2} M_{\sigma,P}(z - qe^{i\theta}) d\theta \right| \\ &\ll \frac{q^4}{(\log N\wp)^2} \ll \left(\frac{1}{N\wp^\sigma} \right)^4 \end{aligned}$$

Note that, by (1) and (3) of Theorem 5.3.4, $M_{\sigma,P}$ is bounded. Thus we see that $M_{\sigma,P}(w)$ converges uniformly to a function, say $M_\sigma(w)$ for $\sigma > 1/2$ (in fact, $1/4$). \square

Remark 5.3.6. We also have, $\int_{\mathbb{C}} M_\sigma(w) |dw| = 1$. But we will show this after showing the next theorem. Note that since, $\int_{\mathbb{C}} M_{\sigma,P}(w) |dw| = 1$ for all P , the uniform convergence already gives,

$$\int_{\mathbb{C}} M_\sigma(w) |dw| \leq 1$$

Theorem 5.3.7. For any $s \in \mathbb{C}$ with $\sigma = \operatorname{Re}(s) > 1$

$$\operatorname{Avg}_\chi \Phi(\mathcal{L}'(\chi, s)) = \int_{\mathbb{C}} M_\sigma(w) \Phi(w) |dw| \quad (5.7)$$

holds for any continuous function Φ of \mathbb{C} .

Proof. From equation 5.2 and Theorem 5.3.4 we have,

$$\begin{aligned} \operatorname{Avg}_\chi (\Phi(\mathcal{L}'_P(\chi, s))) &= \int_{T_P} \Phi(g_{\sigma,P}(t_P)) d^*t_P \\ &= \int_{\mathbb{C}} M_{\sigma,P}(w) \Phi(w) |dw| \end{aligned}$$

The theorem is proved by taking the limit and from Lemma 5.3.3 and Proposition 5.3.5. \square

Note that if we take the particular case of $\Phi(w) = P^{(a,b)}(w) = w^a \bar{w}^b$, Then our results from moments give : (Theorem 4.6.1)

$$(-1)^{a+b} \tilde{\mu}^{(a,b)} = \int_{\mathbb{C}} M_{\sigma}(w) P^{(a,b)}(w) |dw|$$

In particular, taking $a = b = 0$ gives,

$$\int_{\mathbb{C}} M_{\sigma}(w) |dw| = \tilde{\mu}^{(0,0)} = 1$$

Also note that if we consider the Fourier dual of $M_{\sigma}(z)$ given by

$$\tilde{M}_{\sigma}(z) = \int_{\mathbb{C}} M_{\sigma}(w) \psi_z(w) |dw|$$

Then from the above Theorem 5.3.7, we have

$$\tilde{M}_{\sigma}(z) = \text{Avg}_{\chi} \psi_z(\mathcal{L}'(\chi, s))$$

5.4 A NOTE ON HIGHER DERIVATIVES

We note that the above technique theoretically generalizes to higher derivatives. We just need to appropriately choose the $g_{\sigma, \varphi}(t_{\varphi})$ function such that for a local factor, we get

$$\mathcal{L}_{\varphi}^{(n)}(\chi, s) = g_{\sigma, \varphi}(\chi(\varphi)) N_{\varphi}^{-it}$$

But note that for higher derivatives, computing these $M_{\sigma, P}$ functions explicitly becomes very involved. Even in our case we used a computer to simplify the Jacobian.

5.5 POSSIBLE EXTENSION OF OUR RESULT TO $\frac{1}{2} < \sigma \leq 1$

For $\sigma > 1$, the image of $g_{\sigma, P}$ remains bounded as $|P| \rightarrow \infty$. Since the support of $M_{\sigma, P}$ is the image of the mapping, $g_{\sigma, P}$, so the support of M_{σ} is also bounded. Therefore, in the proof of the above theorem we can just assume Φ to be continuous.

This is no longer true for $\sigma > \frac{1}{2}$, i.e. image of $g_{\sigma,P}$ need not be bounded. As remarked earlier, Ihara and Matsumoto, in a later paper [IM11] extends it to $\sigma > 1/2$ under GRH and some conditions on the test function. They introduced the idea of admissible functions and developed a more general notion of $g_{\sigma,\wp}$ which they called g -functions. However their approach does not seem to apply for higher derivatives. It fails at essential steps in section 3.1 and 3.3 of their paper. We are yet to discover a way of doing this and this is currently a work in progress.

BIBLIOGRAPHY

- [BL99] E. Bombieri and J. C. Lagarias, *Complements to Li's criterion for the Riemann hypothesis.*, *Journal of Number Theory*, 77 (1999), 274–287.
- [Bro05] F. C. Brown, *Li's criterion and zero-free regions of L-functions.*, *Journal of Number Theory*, 111 (2005), 1–32.
- [CF76] J.W.S. Cassels and A. Frölich, *Algebraic Number Theory*, Proceedings of an Instructional Conference by London Mathematical Society and International Mathematical Union, Academic press., 1976.
- [CON88] J. B. CONREY, *The fourth moment of derivatives of the Riemann zeta-function*, *The Quarterly Journal of Mathematics*, vol. 39, no. 1, pp. 21–36, doi:10.1093/qmath/39.1.21., March 1988.
- [Dav00] H. Davenport, *Multiplicative Number Theory*, Third edition, Springer Verlag, New York, 2000.
- [Edd18] Sumaia Saad Eddin, *The signs of the Stieltjes constants associated with the Dedekind zeta function*, *Proc. Japan Acad.*, 94, Ser. A, No. 10 (2018).
- [HB10] D.R. Heath-Brown, *Fractional moments of Dirichlet L-functions*, *Acta Arith.*, 145(4), 397–409, 2010.
- [Hec83] Erich Hecke, *Mathematische Werke*, third edition, Vandenhoeck and Ruprecht, Göttingen. With introductory material by B. Schoeneberg, C.L. Siegel, and J. Nielsen. MR 749754 (86a:01049), 1983.
- [HJ77] M. N. Huxley and M. Jutila, *Large values of Dirichlet polynomials IV*, *Acta Arithmetica* 32 (1977), 297–312.

- [HS05] C. Hermite and T. J. Stieltjes, *Correspondance d'Hermite et de Stieltjes, I and II*, edited by B. Baillaud and H. Bourget, Gauthier-Villars, Paris (1905).
- [Iha06] Yasutaka Ihara, *On the Euler-Kronecker constants of global fields and primes with small norms*, Algebraic Geometry and Number Theory, Progr. Math. (Birkhäuser Boston, Boston, MA), Vol. 253, pp. 407–451, 2006.
- [Iha08] Y. Ihara, *On “M-functions” closely related to the distribution of L'/L -values*.
- [IK21] Henryk Iwaniec and Emmanuel Kowalski, *Analytic number theory*, vol. 53, American Mathematical Soc., 2021.
- [IM11] Yasutaka Ihara and Kohji Matsumoto, *On “M-functions” closely related to the distribution of L'/L -values*, Moscow Mathematical Journal, Vol 11, issue 1 (2011), 73–111.
- [IMS09] Y. Ihara, V. Kumar Murty, and Mahoro Shimura, *On the logarithmic derivatives of Dirichlet L-functions at $s = 1$* , Acta Arithmetica, vol 137, 2009.
- [JW35] B. Jessen and A. Wintner, *Distribution Functions and the Riemann Zeta Function*, Trans. Amer. Math. Soc., 38, No. 1, (1935), 48–88.
- [Li97] X.-J. Li, *The positivity of a sequence of numbers and the Riemann hypothesis*, Journal of Number Theory, 65(NT972137) (1997), 325–333.
- [LO77] J. C. Lagarias and A. M. Odlyzko, *Effective versions of the chebotarev density theorem*, Algebraic Number Fields, (A. Fröhlich, Ed.) Proc. of the 1975 Durham Symposium, Academic Press, London & New York (1977), 409–464.
- [Mil11] M.B. Milinovich, *Moments of the Riemann zeta-function at its relative extrema on the critical line*, Bull. Lond. Math. Soc., 43(6), 1119–1129, 2011.
- [MM97] V. Kumar Murty and M. Ram Murty, *Non-vanishing of L-functions and Applications*, Birkhäuser Basel, Springer, 1997.

- [MM15] Mariam Mourtada and Kumar V. Murty, *Distribution of values of $L'/L(\sigma, \chi_D)$* , Moscow Mathematical Journal, Vol 15, issue 3 (2015), 497–509.
- [Mon69] H.L. Montgomery, *Zeros of L-functions.*, Inventiones mathematicae 8 (1969), 346–354.
- [Mon71] ———, *Topics in Multiplicative Number Theory*, Lecture Notes in Math. 227, Springer Verlag, 1971.
- [Mou13] Mariam Mourtada, *The distribution of values of logarithmic derivatives of real L-functions*, PhD Thesis, University of Toronto (2013).
- [Pal31] R. E. A. C. Paley, *On the k -analogues of some theorems in the theory of the Riemann ζ -function*, Proceedings of the London Mathematical Society, Vol s2-32, 1, P.273-311, 1931.
- [Ros00] Michael Rosen, *Number Theory in Function Fields*, Graduate Texts in Mathematics, Springer, 2000.
- [Son14] Keiju Sono, *Upper bounds for the moments of derivatives of Dirichlet L-functions*, Central European Journal of Mathematics, Vol 12(6), 2014.
- [Sou09] K. Soundararajan, *Moments of the Riemann zeta function*, Ann. of Math., 170(2), 981–993, 2009.
- [Sta74] H. M. Stark, *Some effective cases of the Brauer-Siegel Theorem*, Inventiones math. 23, 135-152, 1974.
- [Tit58] E.C. Titchmarsh, *The theory of functions, second ed.*, Oxford University Press, Oxford, 1958.
- [Tsf06] M. A. Tsfasman, *Asymptotic behaviour of the euler-kronecker constant*, Birkhäuser Boston, Boston, MA, 2006.
- [Zha90] W. Zhang, *On the fourth power mean of Dirichlet L-functions*, Chinese Science Bulletin, Vol 35, page 1940-1945, 1990.

COLOPHON

This thesis was typeset using the typographical look-and-feel classicthesis developed by André Miede and Ivo Pletikosić.

The style was inspired by Robert Bringhurst's seminal book on typography "*The Elements of Typographic Style*".